

Common value elections with private information and informative  
priors: theory and experiments

Friederike Mengel and Javier Rivas

No. 44 /15

**BATH ECONOMICS RESEARCH PAPERS**

**Department of Economics**

Department of  
Economics



UNIVERSITY OF  
**BATH**

# Common value elections with private information and informative priors: theory and experiments\*

FRIEDERIKE MENGEL <sup>†</sup>

University of Essex & Maastricht University

JAVIER RIVAS<sup>‡</sup>

University of Bath

February 2, 2016

## Abstract

We study efficiency and information aggregation in common value elections with continuous private signals and informative priors. We show that small elections are not generally efficient and that there are equilibria where some voters vote against their private signal even if it provides useful information and abstention is allowed. This is not the case in large elections, where the fraction of voters who vote against their private signal tends to zero. In an experiment, we then study how informativeness of priors and private signals impact efficiency and information aggregation in small elections. We find that there is a substantial amount of voting against the private signal. Moreover, while most experimental elections are efficient, voting behaviour reveals little about voter's private information.

JEL Classification: D72, D78.

Keywords: Voter turnout, Common Value Elections, Private Information, Swing Voter's Curse, Condorcet Jury Theorem .

---

\*We would like to thank Zia UH Khan, Marina Kuzmenko and Adachi T Madufor for excellent research assistance, Deniz Selman, Fabio Cerina and seminar audiences in Exeter, the European Public Choice Society meeting in Groningen and the Journées LAGCV in Aix-en-Provence for helpful comments. Friederike Mengel thanks the Dutch Science Foundation for financial support (NWO, VENI grant 016.125.040).

<sup>†</sup>Department of Economics, University of Essex, Wivenhoe Park, Colchester CO4 3SQ. *e-mail*: fr.mengel@gmail.com

<sup>‡</sup>Department of Economics, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom. j.rivas@bath.ac.uk, <http://people.bath.ac.uk/fjrr20/>.

# 1 Introduction

Consider a meeting of the executive board of a business where a decision by voting is due on to which of two foreign markets to expand. All members of the executive board have the same target, to increase the profits of the business, yet they may have different opinions about which market will be best for their company. Assume all board members have access to a report detailing which market is likely to be the most profitable one. On top of that, board members may have their own private information based on their past experience, their discussions with other colleagues, etc. The question we ask in this paper is this twofold: can it be rational for board members to ignore their private information and vote following the report even when private information is informative *and* abstention is allowed? Will the committee arrive at the best possible decision given the information they have available?

To answer these questions we consider a common value election between two candidates where voters are not perfectly informed about who is the best candidate. Instead, each voter receives information about the identity of the best candidate from two sources, one public and one private. The public source of information is a common prior shared by all voters. The private source of information consists of an idiosyncratic signal of a certain quality, which could for example reflect the voter's expertise. Each voter knows the quality of his own signal but not the quality of the signals others receive nor these signals themselves. In this setting it may happen that some voters decide to abstain because they believe that their vote is going to harm the chances of the best candidate winning the election. This is known as *strategic abstention* (see for instance McMurray (2010, 2013); Feddersen and Pesendorfer (1996)), which can occur if the signal quality of these voters is low, so that they prefer leaving the decision of selecting a candidate to other, possibly better informed, voters (self-selected experts). In this paper we ask under which conditions a voter may even vote against his signal and what are the implications of such behaviour for efficiency and information aggregation.

From our theoretical analysis we obtain three main results: first, we find that a significant amount of voting against the signal can be observed in equilibrium. Voting against the signal can be rational if the voter deems the signal of too low quality compared to the information contained in the asymmetric (and hence informative) prior. Second, we find that voting does not generally aggregate information efficiently. This is due to a mis-coordination problem that appears as a result of equilibrium multiplicity. Third, for elections with a large number of voters we prove that the effect of an asymmetric common prior vanishes to zero and the election resembles one where the common prior is non-informative.

Our analysis is closely related to McMurray (2013), who studies Condorcet (1785)'s classic common value environment with symmetric priors. The main difference between McMurray

(2013) and the present paper is that we allow for the common prior to be asymmetric: i.e. not all candidates are equally likely to be the best one a priori. This gives rise to a phenomenon not present in McMurray (2013): voters can vote against their own signal. With symmetric priors any signal is at least as good as the prior in predicting the best candidate. This means that no voter has incentives to vote against his signal and their decision then reduces to whether to abstain or not. In our paper the fact that a signal may be less informative than the common prior means that some voters will choose to vote against their private signal. Contrary to previous literature, voting against the private signal is not the result of the different biases voters may have (Feddersen and Pesendorfer, 1996; Rivas and Rodríguez-Álvarez, 2012).

Our experiments test both the predictions of the symmetric case studied in McMurray (2013) as well as the predictions of the asymmetric case introduced here for small elections. As expected from the theoretical analysis, few voters ( $< 10\%$ ) vote against their signal with uninformative (symmetric) priors, but 40 – 80%, depending on signal accuracy, do so in the case with informative (asymmetric) priors. Turnout is higher in the asymmetric case (83 – 86%) than in the symmetric case (78%) and slightly higher than theoretically expected. The experiments deliver a surprising result in terms of efficiency. While as expected more informative priors lead to higher efficiency, more informative signals do not. This is because in the experiments voters trust moderately precise signals too much, which leads to an inefficient use of information. A consequence of this, efficiency is higher with less informative signals.

Our research contributes to the literature on common value elections and strategic abstention. The classic paper of Austen-Smith and Banks (1996) raised serious questions about Condorcet’s implicit assumption that all voters will vote naively, i.e. vote as if they were the only voter. They showed that voting against the signal can arise if abstention is not allowed and all voters have the same signal quality. In Feddersen and Pesendorfer (1996) voters are of three types: partisans, fully informed and uninformed. Partisans support a certain candidate irrespective of the information available while fully informed and uninformed voters prefer the best candidate. Fully informed voters know for certain who is the best candidate while uninformed voters have no information about the best candidate other than the common prior. They show that a positive fraction of uninformed voters abstain even when they strictly prefer one candidate over the other (swingers voter’s curse). Battaglini et al. (2010) experimentally tested this model and found results in terms of efficiency, turnout and the margin of victory that are in line with theory. We find theoretically and experimentally that being uninformed is not a requirement for the swingers voter’s curse (see also McMurray (2013)). Indeed, the fact that voters possess information of different qualities leads to a self selection in abstention; those with lower quality signals abstain, even if their signal is more

informative than the prior, and even if based on the information they have they strictly prefer one candidate over the other.

In Feddersen and Pesendorfer (1997) voters receive information from different sources, where each source may provide information of different qualities. However, they do not allow for abstention, which is a crucial difference to our model. Feddersen and Pesendorfer (1998) allow voters to abstain. However, all voters receive information of the same quality. The results in Feddersen and Pesendorfer (1998) are similar to ours except for the fact that in their article the reason behind what each voter chooses given his signal is how biased towards either candidate he is. In our paper, no voter is biased and the driving force behind what each voter chooses given his signal is the quality of the signal. Our paper hence identifies a new mechanism of why voting behaviour can differ across players: differences in behavior might be due either to differences in preferences, as in Feddersen and Pesendorfer (1998), or due to differences in the perceived quality of the information, as in our case.

Also related are Ben-Yashar and Milchtaich (2007) who study voters with homogeneous preferences and private signals of different qualities. However, they do not consider the possibility of abstention; their focus is on computing the best monotone voting rule. Krishna and Morgan (2012) investigate the welfare effects of introducing voluntary voting when all voters have the same signal quality. Oliveros (2013) presents a model where voters can buy information of different qualities and studies the effects of different ideologies on information acquisition.

A technical difference between our paper and some of the previous theoretical literature (McMurray (2013), Feddersen and Pesendorfer (1996, 1997, 1999) among others) is that we do not consider an uncertain number of voters, i.e. Poisson games (Myerson, 1998), to prove our results. In elections with a small number of voters, as it is the case in the example in our starting paragraph, this assumption may seem hard to justify. The fact that we do not consider Poisson games does not lead to different results when the number of voters is large. In this case, our results mirror those of McMurray (2013).

Finally, our research also contributes to the experimental literature on the Condorcet Jury paradigm. The first published experiment on behaviour in voting games was conducted by Guarnaschelli et al. (2000) who base their experiment on Feddersen and Pesendorfer (1998)'s analysis of strategic voting with the unanimity rule. Closer to our setting are Battaglini et al. (2008) and Battaglini et al. (2010) as described above, and also Morton and Tyran (2011) who extend the Battaglini et al. (2008) setting by exploring an environment where poorly informed voters are not completely uninformed - they simply receive lower quality signals. This can lead to equilibria where all voters vote and to equilibria where the poorly informed voters abstain. However, as priors are symmetric, there should not be any voting against the

signal in this setting.

Studying experimentally whether participants would be willing to vote against their signal when it is rational to do so raises interesting questions in itself. Violations of Bayesian updating are widely documented in experimental research and there are two types of biases which would lead to opposite results in terms of participants voting against their signal. The well documented phenomenon of base-rate neglect (Kahnemann and Tversky, 1972; Grether, 1980; Erev et al., 2008) will lead agents to overweight sample information and hence would imply that voting against the signal is not commonly observed in the experiment. However, there is also the opposing phenomenon of conservatism (Ward, 1982) which implies that participants overweight the prior and hence would reinforce the strategic incentives to vote against the signal. By studying for the first time common value elections with informative priors our experiment can shed some light on the role of these two opposing biases in strategic voting.

The rest of the paper is organized as follows. In section 2 we introduce the model and present the main theoretical results. In section 3 we describe the design of the experiments and present the experimental results. In section 4 present further theoretical results where we consider the limit case then the number of voters grows large. Finally, section 5 concludes.

## 2 Theory

### 2.1 The Model

Consider a setting where  $N + 1 \geq 2$  voters have to decide between candidate  $R$ (ed) or candidate  $B$ (lue) by simultaneously casting a vote for either candidate or abstaining. The candidate that the receives the most votes wins the election. In case of a tie each candidate wins with equal probability.

Each voter derives one unit of utility if the candidate who wins coincides with the state of nature and zero units of utility otherwise. The state of nature is a random variable  $s \in \{R, B\}$  where without loss of generality we assume that the probability that the state is  $R$  is given by  $p \geq \frac{1}{2}$ .<sup>1</sup> We restrict our attention to situations where  $p \in [\frac{1}{2}, 1)$  as if  $p = 1$  then all voters agree that  $R$  is the best candidate and thus will vote for him regardless on any other information they may have available. The value of  $p$  is common knowledge and we refer to it as the common prior.

---

<sup>1</sup>The assumption  $p \geq \frac{1}{2}$  is without loss of generality as if  $p < \frac{1}{2}$  then a relabeling of  $R$  to  $B$  and vice-versa makes the analysis that follows still valid.

Before the election, each voter  $i$  receives a signal  $\sigma_i \in \{R, B\}$  with quality  $q_i \in [\frac{1}{2}, 1]$  where

$$P(\sigma_i = s | q_i) = q_i.$$

Given the state of nature, signals of different voters are conditional independent. Both the signal received by each voter as well as the quality of such signal are private information. The distribution of signal qualities for each voter in the population is common knowledge, identical, independently distributed and given by the strictly increasing cumulative density function  $F : [\frac{1}{2}, 1] \rightarrow [0, 1]$  and integrable probability density function  $f : [\frac{1}{2}, 1] \rightarrow \mathbb{R}^+$ . Define the average signal quality as  $\mu = \int_{\frac{1}{2}}^1 q f(q) dq$  and consider  $\mu \in (\frac{1}{2}, 1)$  to avoid the trivial cases where all voters receive a useless signal or when all voters receive a perfectly informative signal.

Thus, before the election each voter knows the common prior, his own signal and the quality of such signal, as well as the distribution of the quality of other voters' signals. However, he ignores the state of nature, the signals received by other voters, and the quality of such signals.

A strategy for each voter is a map  $v : \{R, B\} \times [\frac{1}{2}, 1] \rightarrow \{\emptyset, R, B\}$  where  $v(\sigma_i, q_i)$  is the action of voter  $i$  who receives signal  $\sigma_i$  of quality  $q_i$ , and  $\emptyset$  stands for the action of abstaining. Note that we focus on symmetric strategies: voters that are the same (same signal and quality) behave the same. The fact that we only consider symmetric equilibria does not undermine our main findings: if voting against the signal is possible in an equilibrium with symmetric strategies then it is also possible in an equilibrium when asymmetric strategies are considered. Moreover, as we argue later on in section 4.1, when there is a large number of voters considering symmetric strategies is without loss of generality.

Note that unlike most recent papers on voting and information sharing we do not assume a Poisson distribution for the number of voters (see the seminal work by Myerson (1998) and Myerson (2000) and more recent references by Myatt (2012) and Nunez (2010) among others). This assumption is often employed given its technical conveniences, namely, independent common public information and independence of actions. However, a drawback of assuming a Poisson distribution for the number of voters is that voters are uncertain of how many other voters there are in the population. While this seems a suitable assumption in large elections, with small elections (committees, for example), which are the focus of this paper, it seems unreasonable to assume that voters ignore how many other voters there are.

Given that we focus on a small number of voters, one may be interested in knowing how allowing voters to deliberate would affect our setting. However, since we are dealing with a common value election, the addition of a deliberation stage in the form of a straw poll a la

Coughlan (2000) does not affect the strategic incentives of voters and, hence, will not affect our results.

## 2.2 Analysis

When a voter decides whether to vote for  $R$ ,  $B$  or to abstain, he compares the payoff he obtains under these three actions given the actions of all other voters. However, a voter can influence his own payoff only when his vote can change the outcome of the election (i.e. he is pivotal). This can happen if and only if candidates  $R$  and  $B$  are at most one vote apart when counting the votes of the other  $N$  voters. Thus, let  $\pi_t(v, s)$  be the probability that candidate  $R$  receives the same number of votes as candidate  $B$  (i.e. there is tie) when  $N$  voters use strategy  $v$  and the state is  $s$ . Similarly, let  $\pi_R(v, s)$  be the probability candidate  $R$  receives exactly one vote less than candidate  $B$  when  $N$  voters use strategy  $v$  and the state is  $s$ . Finally, let  $\pi_B(v, s)$  be the probability candidate  $B$  receives exactly one vote less than candidate  $R$  when  $N$  voters use strategy  $v$  and the state is  $s$ .

Before we write down the payoff each voter obtains from playing the three different actions, it is useful to understand how likely each state is when a voter only considers his available information (i.e. ignoring strategic considerations). We have the following:

$$\begin{aligned} P(s = R | \sigma_i = R, q_i) &= \frac{pq_i}{pq_i + (1-p)(1-q_i)}, \\ P(s = B | \sigma_i = R, q_i) &= \frac{(1-p)(1-q_i)}{pq_i + (1-p)(1-q_i)}, \\ P(s = R | \sigma_i = B, q_i) &= \frac{p(1-q_i)}{p(1-q_i) + (1-p)q_i}, \\ P(s = B | \sigma_i = B, q_i) &= \frac{(1-p)q_i}{p(1-q_i) + (1-p)q_i}. \end{aligned}$$

Notice that the private signal of voter  $i$  is more informative than the prior,  $P(s | \sigma_i = s, q_i) \geq \frac{1}{2}$  for all  $s \in \{R, B\}$ , if and only if  $q_i \geq p$ .

The expected utility voter  $i$  derives from voting for  $R$  compared to voting for  $B$  when the other  $N$  voters use strategy  $v$  is then given by

$$\begin{aligned} u_i(R, B, v) &= P(s = R | \sigma_i, q_i) \left[ \pi_t(v, R) + \frac{1}{2}\pi_B(v, R) + \frac{1}{2}\pi_R(v, R) \right] \\ &\quad - P(s = B | \sigma_i, q_i) \left[ \pi_t(v, B) + \frac{1}{2}\pi_B(v, B) + \frac{1}{2}\pi_R(v, B) \right]. \end{aligned} \quad (1)$$

In words, if the state is  $R$  then the increase in payoff from voting  $R$  instead of  $B$  is: 1 if there is a tie when counting all other  $N$  votes (the best candidate wins),  $\frac{1}{2}$  if  $B$  is one vote behind (the best candidate is chosen as opposed to forcing a tie), and  $\frac{1}{2}$  if  $R$  is one vote



behind (a tie is forced as opposed to not having the best candidate win). On the other hand, if the state is  $B$  then the increase in payoff from voting  $R$  instead of  $B$  is:  $-1$  if there is a tie when counting all other  $N$  votes (the best candidate does not win),  $-\frac{1}{2}$  if  $B$  is one vote behind (the best candidate is not chosen as opposed to forcing a tie), and  $-\frac{1}{2}$  if  $R$  is one vote behind (a tie is forced as opposed to having the best candidate win).

Similarly, the expected utility voter  $i$  derives from voting for  $R$  or  $B$  compared to abstaining when the other  $N$  voters use strategy  $v$  is given respectively by

$$u_i(R, \emptyset, v) = P(s = R | \sigma_i, q_i) \left[ \frac{1}{2} \pi_t(v, R) + \frac{1}{2} \pi_R(v, R) \right] - P(s = B | \sigma_i, q_i) \left[ \frac{1}{2} \pi_t(v, B) + \frac{1}{2} \pi_R(v, B) \right], \quad (2)$$

$$u_i(B, \emptyset, v) = P(s = B | \sigma_i, q_i) \left[ \frac{1}{2} \pi_t(v, B) + \frac{1}{2} \pi_B(v, B) \right] - P(s = R | \sigma_i, q_i) \left[ \frac{1}{2} \pi_t(v, R) + \frac{1}{2} \pi_B(v, R) \right]. \quad (3)$$

To simplify the exposition, we assume that if voters are indifferent between the two candidates they prefer the one that coincides with their signal. Similarly, if voters are indifferent between voting for a certain candidate or abstaining, they follow their signal. As it will be clear later on, the fact that  $f$  is integrable means that the probability that a voter is indifferent between two options (voting to one candidate or the other, or voting to either candidate or abstaining) is zero. As such, the way indifference ties are broken has no effect in our results and it also allows us to ignore mixed strategies.

A voter who receives signal  $R$  votes for  $R$  if and only if  $u_i(R, B, v) \geq 0$  and  $u_i(R, \emptyset, v) \geq 0$ . Such voter abstains if and only if  $u_i(R, \emptyset, v) < 0$  and  $u_i(B, \emptyset, v) \leq 0$ , and votes for  $B$  if and only if  $u_i(R, B, v) < 0$  and  $u_i(B, \emptyset, v) > 0$ . Thus, expressions (1), (2) and (3) are what determines how a voter behaves given how the other voters behave.

We have the following characterization of all symmetric equilibria (all mathematical proofs are presented in the appendix):

**Theorem 1.** *There exists an equilibrium. The equilibrium is either of two types:*

- *Type 1, characterized by two cutpoints  $\frac{1}{2} \leq q_B^- \leq q_B^+ \leq 1$  with  $q_B^- \leq p$  such that*

$$v(\sigma_i, q_i) = \begin{cases} R & \text{if either } \sigma_i = R \text{ or } \sigma_i = B \text{ and } q_i < q_B^-, \\ B & \text{if } \sigma_i = B \text{ and } q_i \geq q_B^+, \\ \emptyset & \text{otherwise.} \end{cases}$$

- Type 2, characterized by two cutpoints  $\frac{1}{2} \leq q_R^+ \leq q_B^+ \leq 1$  such that

$$v(\sigma_i, q_i) = \begin{cases} R & \text{if } \sigma_i = R \text{ and } q_i \geq q_R^+, \\ B & \text{if } \sigma_i = B \text{ and } q_i \geq q_B^+, \\ \emptyset & \text{otherwise.} \end{cases}$$

In equilibrium of Type 1 all voters who receive signal  $R$  vote and they do so for candidate  $R$ . These are the voters who receive a signal that agrees with the common prior. On the other hand, voters who receive a signal against the common prior, i.e. signal  $B$ , behave as follows: those with a low quality signal ignore their signal and vote according to the common prior, those with a moderately informative signal abstain, and those with a sufficiently informative signal vote according to their signal.

In equilibrium of Type 2 no voter votes against his signal. Note that  $q_R^+ \leq q_B^+$  implies that those voters who receive a signal that agrees with the common prior are less likely to abstain than those who receive a signal against. This is the case because  $p \geq \frac{1}{2}$  and, thus, if a voter receives signal  $R$  the common prior makes him trust his signal more whereas if voter receives signal  $B$  he is less convinced about candidate  $B$  than his signal quality suggests as the common prior goes against  $B$ .

The reason why there is not an equilibrium where voters who receive signal  $R$  vote for  $B$  is that  $p \geq \frac{1}{2}$  and, thus, a voter whose signal agrees with the common prior believes that  $R$  is the best candidate so he either abstains or votes for  $R$ . Figures 1 and 2 present a graphical representation of both types of equilibria.

Figure 1: Equilibrium of Type 1





Figure 2: Equilibrium of Type 2



The expected fraction of voters who vote against their signal in equilibrium of Type 1 is given by  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq$  which, as we shall see with examples, can be a strictly positive number. The fraction of voters who abstain is given by  $\int_{q_B^-}^{q_R^+} f(q) dq$  in equilibrium of Type 1 and  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq + \int_{\frac{1}{2}}^{q_R^+} f(q) dq$  in equilibrium of Type 2. This is the so-called strategic abstention (and swing voters curse), found for instance in McMurray (2013) and Feddersen and Pesendorfer (1996) (see McMurray (2010) for an empirical reference).

The reason why some voters vote against their signal is the following. Consider a very simple example where there are only two voters. In this case a voter is always pivotal and thus learns very little from the fact that he is pivotal (he still does learn some information, as there are three different possibilities for a voter to be pivotal). In this case if a voter receives a low quality signal against the common prior, given that he does not learn much from being pivotal, he may still prefer to vote for what the common prior suggests if the common prior is informative enough. This reasoning extends to more than just two voters. Assume that a voter receives a low quality signal supporting the candidate that goes against the common prior. If such voter is pivotal, he knows that there are mixed signals in the population, which suggests that the common prior may be wrong. However, his information may still support the same candidate as the common prior given that his updated belief stills put a significant probability on such candidate because the voter's signal is of low quality. Thus, the voter may have incentives to disobey his signal and vote against it.

The reason why strategic abstention is possible is that if a voter receives a signal of moderate quality and the common prior is not very informative (or he receives a signal of high quality against an informative the common prior, but not of sufficiently high quality), then if the voter is pivotal he may prefer to abstain and leave the decision to those who are presumably better informed. This is because if the voter is pivotal there is a significant chance that the best candidate is ahead by one vote as opposed to the other candidate ahead by one vote or there being a tie. Hence, by voting the voter runs the risk of contradicting

the opinion of most other voters who do not abstain and who have a better signal quality than himself. In this situation the voter is better off by abstaining, even if he prefers one candidate over the other, and leaving the decision of electing a candidate to the other more informative voters.

Note that from the information revelation point, voting against the voter's signal is worse than abstaining. When a voter abstains he reveals that his signal is not very informative. However, in an equilibrium where voters may vote against their signal, if a voter votes for  $R$  it is not clear whether such voter received signal  $R$  or  $B$ . That is, voting against the signal harms the chances of the best candidate winning the election more than abstention.

Theorem 1 states that in an equilibrium of Type 1,  $q_B^- \leq p$ . Numerical examples shows that this inequality can be strict. If instead of a group of voters a single voter (dictator) chose the winning candidate, straightforward calculations show that this voter will choose to follow his signal if and only if his signal points at candidate  $R$  or if it points at candidate  $B$  and the signal quality is at least  $p$ . In the language of the model, if  $N + 1 = 1$  then the unique equilibrium is Type 1 with  $q_B^- = q_B^+ = p$ . Thus, the fact that the group of voters includes more than just one voter means that voters are less likely to vote against their signal. That is, more voters means that each of them has more incentives to share their signal even if such signal is of a quality lower than the prior. Later in the paper we show that the fraction of voters who vote against their signal converges to zero as the number of voters increases to infinity.

As discussed in the introduction, McMurray (2013) considers a setting very similar to ours where the main difference is that he assumes  $p = \frac{1}{2}$ . The consequence of this is that in his setting the only possible equilibria is Type 2 with  $q_R^+ = q_B^+$ . The fact that  $p > \frac{1}{2}$  is what allows the existence of an equilibrium of Type 1 with  $q_B^- > \frac{1}{2}$  and an equilibrium of Type 2 with  $q_R^+ < q_B^+$ . The comparison of our results to McMurray (2013) is explored in more detail later on when we consider elections with a large number of voters.

It is worth pointing out the similarities between our result in Theorem 1 and Proposition 1 in Feddersen and Pesendorfer (1998) (particularly striking is the resemblance between figures 1 and 2 and figure 1 in Feddersen and Pesendorfer (1998)). However, both results originate from very different sources. In our paper, voters' behavior depends on the signal they receive, but also on the quality of such signal. In Feddersen and Pesendorfer (1998), voters' behavior depends on the signal they receive and on their bias towards each of the candidates. Thus, the fact that unbiased voters receive signals of different qualities mimics the behavior observed when biased voters receive information of equal quality. A fundamental difference between these two situations is that a voter who is biased takes such bias as given while an unbiased voter is aware of the fact that his signal may or may not be very accurate.

### 2.2.1 Numerical Examples

Next we present some examples that illustrate the results of Theorem 1. In tables 1 and 2 we calculate some of the possible equilibria when there are 4 and 5 voters respectively and signal qualities are distributed uniformly. The parameter constellations in table 1 are the ones used in the experiments.

Table 1: Equilibria,  $N + 1 = 4$

$p = 0.5, q \sim U [\frac{1}{2}, 1]$	$p = 0.95, q \sim U [\frac{1}{2}, 1]$	$p = 0.95, q \sim U [\frac{1}{2}, \frac{3}{4}]$
$q_R^+ = 0.67$	$q_B^- = 0.54$	$q_B^- = 0.57$
$q_B^+ = 0.67$	$q_B^+ = 0.86$	$q_B^+ = 0.76$

Table 2: Equilibria,  $N + 1 = 5$

$p = 0.5, q \sim U [\frac{1}{2}, 1]$		$p = 0.8, q \sim U [\frac{1}{2}, 1]$		$p = 0.95, q \sim U [\frac{1}{2}, 1]$	
$q_R^+ = 0.5$	$q_R^+ = 0.68$	$q_R^+ = 0.60$	$q_B^- = 0.59$	$q_R^+ = 0.51$	$q_B^- = 0.68$
$q_B^+ = 0.5$	$q_B^+ = 0.68$	$q_B^+ = 0.75$	$q_B^+ = 0.59$	$q_B^+ = 0.82$	$q_B^+ = 0.68$

Numerical results suggest that the equilibrium is unique if and only if  $N + 1$  is even and that if  $N + 1$  is odd then there are exactly two equilibria. However, we have been unable to prove this formally. The problem of uniqueness of equilibrium in voting models such as this far from trivial (McMurray (2013)) and is often ignored (Feddersen and Pesendorfer (1996, 1997, 1999)). Nevertheless, uniqueness of equilibrium is not necessary for our results. Our characterization in Theorem 1 together with the examples above already illustrate one of the points of this paper: the possibility for rational voters voting against their signal when such signal is informative and abstention is allowed. On top of that, uniqueness of equilibrium is also not required for the experimental results; we are not interested in making point-wise prediction but in understanding whether the behavior of subjects responds the way the theoretical results predict when we change the parameters of the model.

With respect to whether or not information is used efficiently in equilibrium, the examples in Table 2 already tell us that this is not necessarily the case. For instance, if  $N + 1 = 5$  and  $p = 0.8$  then the probability with which the best candidate wins is 0.92 in Equilibrium of Type 1 and 0.93 in the equilibrium of Type 2. Numerical results show that this difference tends to increase if there are fewer voters. However, as we shall show in the section 4.1, for elections with a large number of voters the best candidate wins the election with probability one.

### 2.3 Testable Predictions and Empirical Questions

In this section we describe some key qualitative properties that we are interested in testing in the laboratory. One of the new findings in this paper is that, unlike in the symmetric prior case studied in McMurray (2013), participants may vote against their signal if it is not accurate enough. As discussed above, this can have important consequences for information aggregation. We ask the following question:

**Voting against one’s signal** *How is the propensity to vote against the signal affected by changes in the prior  $p$  and the signal accuracy  $q$ ?*

From theory we would expect to see voting against the prior only if priors are asymmetric and if signal accuracy is “low enough”. Empirically, whether this prediction holds depends crucially on how people update their prior on the basis of the information they received. Two different failures of Bayesian updating have been robustly documented in the literature: (i) base-rate neglect, which leads to overweighting sampled information (Kahnemann and Tversky (1972); Grether (1980); Erev et al. (2008)) and (ii) conservatism, which leads to underweighting or even ignoring the sample (Ward (1982)).

Base rate neglect is not important if priors are symmetric ( $p = 0.5$ ). With asymmetric priors, however, it could potentially play an important role. Under base-rate neglect participants would vote with their signal more often, leading to more information being revealed, but possibly worse outcomes in terms of the efficiency of the majority decision. Conservatism would lead to the opposite prediction. Participants would vote with the prior too often leading to both worse information aggregation and lower efficiency. Hence, while theory might be a good predictor of behavior for symmetric priors, its predictive accuracy could be far worse in the case of asymmetric priors if base-rate neglect or conservatism play important roles in this setting. If participants vote too often or too seldom against their signal, information revelation and efficiency are impacted as well. We hence ask:

**Information Revelation** *How is information revelation affected by changes in the prior  $p$  and the signal accuracy  $q$ ?*

**Efficiency** *How is the efficiency of voting outcomes affected by changes in the prior  $p$  and the signal accuracy  $q$ ?*

We would expect efficiency to increase both as priors become more asymmetric (hence containing more information) and as signals become more accurate. However, in the presence

of biases, such as base-rate neglect or conservatism, this may not necessarily be the case. Our experiments will provide an empirical test of how the symmetric and asymmetric settings differ with regard to these issues and how potential biases affect the explanatory power of the theory in both these two settings.

## 3 Experiments

### 3.1 Design of the Experiments

Our experiment implements the setting described in the theoretical section for  $N + 1 = 4$ , i.e. four voters. In all treatments participants played a voting game for 30 rounds. After each round they were randomly re-matched in a new group of four voters. Each round proceeded as follows. First, participants were reminded of the value of  $p$ . They were then shown their private signal and informed about the accuracy of their signal  $q_i$ . They were afterwards asked to vote for either RED, BLUE or to ABSTAIN, where the order of the first two options was randomized. At the end of each round they were informed about their own vote, the majority vote in the group, the realized state and their payoff. Participants received 10 experimental tokens if the majority vote matched the state and 2 tokens if it did not. At the end of the experiment one round was randomly drawn and participants were paid for that round only plus a show up fee of 3 tokens. Tokens were converted into GBP at a rate of 1:1.

To answer our questions regarding information revelation and efficiency we systematically vary  $p$  and  $q$ . Treatment **SYM** implements the symmetric setting analyzed by McMurray (2013). Both states are equally likely and signal accuracy is distributed uniformly in  $[0.5, 1]$ . In treatment **ASYM** an asymmetric prior of 0.95 is implemented. Treatment **ASYM-COARSE** coincides with treatment **ASYM**, but the signal accuracy is lower:  $q \sim U[0.5, 0.75]$ . In each treatment we had 24 participants organized in three matching groups (clusters) of size 8. Theoretical predictions for these different cases can be found in Section 2.2.1 Table 1.

The experiments were conducted in May 2015 at EssexLab at the University of Essex. Participants earned either 13 GBP or 5 GBP depending on whether, in the round randomly drawn for payment, the majority vote matched the state or not.<sup>2</sup> The experiment lasted around 45min, it was programmed in z-tree (Fischbacher, 2007) and participants were recruited using hroot.

---

<sup>2</sup>In May 2015, 13 GBP equalled about 20.50 US dollars and 5 GBP around 7.90 US dollars.

## 3.2 Experimental Results

We first look at overall outcomes in terms of efficiency and information revelation and then look at the individual behaviour underlying these aggregate effects.

### 3.2.1 Aggregate Outcomes

We start with the efficiency of voting outcomes. We would expect efficiency to increase both as priors become more asymmetric (hence containing more information) and as signals become more accurate. In terms of our treatments we would hence expect higher efficiency in **ASYM** compared to **SYM** and higher efficiency in **ASYM** compared to **ASYM-COARSE**.




Figure 3: Efficiency: Panel (a) shows the percentage of time the majority vote agreed with the state over time across the three treatments. Panel (b) shows the percentage of times participants vote with their signal over time across the three treatments.

Figure 3 (Panel (a)) shows efficiency over time across the three treatments. As expected, efficiency is higher in **ASYM** compared to **SYM**, even though they are converging over time and across the last 5 periods they are no longer statistically different (Table 3). By contrast, efficiency is higher in **ASYM-COARSE** compared to **ASYM**, which is not what we expected. The difference is not statistically significant across the first five periods (column (1) in Table 3), but it is significant across all rounds (column (2)) and significant and substantial across the last five rounds (column (3)). In fact, in **ASYM-COARSE** the vote is almost always efficient. We will explore the reasons for this effect below. There are also some interesting time trends. While in **ASYM** efficiency decreases over time (OLS coefficient  $-0.0039^{***}$ ), in **SYM** efficiency increases over time ( $0.0064^{***}$ ) stabilizing at around 80 percent efficient votes.<sup>3</sup>

<sup>3</sup>OLS coefficients in brackets are from random effects OLS regressions (including a constant) where the efficiency measure is regressed on period. Standard errors are clustered by matching group.



VARIABLES	(1) All periods	(2) periods 1-5	(3) periods 26-30
<b>ASYM</b>	0.169*** (0.0160)	0.317*** (0.0434)	0.0417 (0.0440)
<b>ASYM-COARSE</b>	0.231*** (0.0160)	0.325*** (0.0434)	0.183*** (0.0440)
Constant	0.751*** (0.0113)	0.625*** (0.0307)	0.783*** (0.0311)
Observations	2,160	360	360
Participants	72	72	72

Robust standard errors in parentheses

\*\*\* p<0.01, \*\* p<0.05, \* p<0.1

Table 3: Random effects OLS regression of efficiency on treatment dummies. Note: (\*\*\*) significant at 1% level, (\*\*) at 5% level and (\*) at 10% level. Standard errors are clustered by matching group.

**Result 1** *Voting outcomes are most efficient in **ASYM-COARSE** followed by **ASYM** and **SYM**.*

**Information Revelation** Next, we study information revelation, i.e. how much about participants' signals (and hence the state) is revealed via their voting behavior. Panel (b) in Figure 3 shows that the share of participants who vote with their signal is roughly constant across time and about the same across treatments. This, however, does not tell us much about how much information is revealed since it is important to understand who votes against their signal, i.e. is it participants whose signal coincides with the prior or those whose signal differs from the prior? and how accurate is the signal of those who vote against it?

One can either study information revelation by making inference based on the theoretical benchmark or in a purely empirical manner. We will discuss both in turn. As mentioned above, in all the equilibria in **ASYM** and **ASYM-COARSE** only participants with a signal that goes against the prior would ever abstain. Hence, if we make inference based on the theoretical benchmark, the decision to abstain reveals the signal as does the decision to vote against the prior. The decision to vote with the prior, by contrast, does not reveal (in a deterministic sense) the signal. Can we make inference on votes that follow the prior? The problem with such inference is that it depends on assumptions about specific equilibria being played. In particular, the conditional probability  $\Pr(\sigma|v_i = BLUE)$  will vary depending on the equilibrium under question. Hence, any theoretical inference be conditional on particular equilibria. For this reason we focus on a purely empirical measure of information revelation.

In particular, we use the ratio of post- to pre-voting Shannon entropy:

$$IR = \frac{-\sum_{s \in \{R,B\}} \hat{p}(s) \log \hat{p}(s)}{-(p \log p + (1-p) \log(1-p))},$$

where  $\hat{p}(s)$  denotes an outside observer's belief on the state after observing the distribution of votes (but without knowledge of the signals received by voters). In particular  $\hat{p}(s)$  denotes the probability that an outside observer (endowed with an uninformative prior), who knows the empirical frequencies  $\Pr(\sigma_i, q_i | v_i)$ , attaches to state  $s$  after seeing the vote. If  $IR < 1$ , then this outside observer will be more informed about  $s$  compared to someone who just knows the prior and if  $IR > 1$ , then the prior reveals more information about  $s$  than the vote.

	RED state			BLUE state		
	Votes	$\sigma = RED$		Votes	$\sigma = BLUE$	
<b>SYM</b>	RED	0.57	0.95 (0.76)	RED	0.23	0.18 (0.63)
	BLUE	0.15	0.26 (0.70)	BLUE	0.57	0.86 (0.74)
	abstain	0.28	0.57 (0.59)	abstain	0.20	0.45 (0.59)
<b>ASYM</b>	RED	0.66	1 (0.85)	RED	0.23	0.20 (0.67)
	BLUE	0.33	0.5 (0.6)	BLUE	0.57	0.80 (0.74)
	abstain	-	-	abstain	0.20	0.33 (0.62)
<b>ASYM-COARSE</b>	RED	0.40	0.85 (0.61)	RED	0.04	0.52 (0.63)
	BLUE	0.40	0.60 (0.61)	BLUE	0.81	0.79 (0.62)
	abstain	0.20	0.60 (0.59)	abstain	0.15	0.57 (0.61)

Table 4: Vote Distribution and share of participants voting (RED, BLUE or abstaining) overall and conditional on observing a RED signal (in brackets average signal accuracy) for each of the three treatments and the two states.

Table 4 gives an overview of voting behaviour. In **SYM**, participants tend to vote RED or BLUE with about equal frequencies. However, there is also a substantial share of abstentions. Consistently with theory, voters rarely vote against their signal in this condition. Since the setting is symmetric, there are, as expected, no differences between RED and BLUE signals. Let us compute the population posterior for RED, i.e.  $\hat{p}(RED)$ . In the red state, observing someone vote RED is indicative of a signal RED with probability 0.95 with average accuracy of 0.76, observing an abstention reveals RED with probability 0.57 (average accuracy 0.59) and observing BLUE indicates RED with probability 0.26 (average accuracy 0.70). The population posterior  $\hat{p}(RED)$  is hence 0.533 in the red state and 0.46 in the blue state, implying  $IR_{\text{SYM}} < 1$ . Hence knowledge of the empirical frequencies  $\Pr(\sigma_i, q_i | v_i)$  as well as the vote distribution is informative about the state, as in the RED state  $\hat{p}(RED) > 0.5$  and in the BLUE state  $\hat{p}(RED) < 0.5$ . The population posterior reveals marginally more information about the state than the uninformative prior.

In treatment **ASYM** most voters vote with BLUE, but there is also a substantial share of abstention and a significant share of voting for RED. Observing someone vote RED is indicative of a signal RED with probability 1 with average accuracy of 0.85 while observing BLUE indicates RED with probability 0.33 (average accuracy 0.6). Two problems with information revelation become apparent. First, since as predicted by theory a substantial share of participants who vote for BLUE actually saw a RED signal, it is harder to make inference from observing BLUE votes. Second, and this goes against theoretical insights, 15 percent of those voting for RED did actually see a BLUE signal. The population posterior  $\hat{p}(RED)$  is 0.76 in the red state and  $\hat{p}(RED) = 0.41$  in the blue state implying  $IR_{ASYM} > 1$ . The population posterior reveals less information compared to the prior, but both are informative in this case.<sup>4</sup> In treatment **ASYM-COARSE** patterns are similar to those in **ASYM**, but lower signal quality means worse inference compared to treatment **ASYM**, but still information is lost via voting. The population posterior is  $\hat{p}(RED) = 0.61$  in the red state and  $\hat{p}(RED) = 0.46$  in the blue state

As we expected from theory, the setting with asymmetric priors is problematic in terms of information revelation. An outside observer, who knows the empirical frequencies  $\Pr(\sigma_i, q_i | v_i)$  as well as the vote distribution, will have a less accurate idea of the state  $s$  compared to someone just relying on the prior in both treatments **ASYM** and **ASYM-COARSE**. It should be noted, though, that in all treatments voting is informative with  $\hat{p}(RED) > 0.5$  in the RED state and  $\hat{p}(RED) < 0.5$  in the BLUE state.

**Result 2** *Voting outcomes are informative in all treatments. Voting reveals more information than the (uninformative) prior in **SYM**, but less information than the prior in **ASYM** and **ASYM-COARSE**.*


### 3.2.2 Individual Behaviour

In this subsection we turn our attention to individual voting strategies to get a better understanding for what underlies our aggregate findings.

Figure 4 illustrates how voters with RED (left panel) and BLUE (right panel) signals vote in treatment **SYM**. As expected, we don't see substantial differences between the two cases. Irrespective of the signal received, only very few participants vote against their signal. The share of abstentions is high (around 60%) if the signal is uninformative and decreases sharply around  $q \approx 0.6$  in line with theoretical predictions. The fact that voters largely vote

---

<sup>4</sup>Note that there are only few instances where the state is RED in treatments **ASYM** and **ASYM-COARSE**. If we aggregate across both states results are very similar.




(a) RED signal

(b) BLUE signal

Figure 4: The figure shows the vote distribution (red, abstain, blue) conditional on signal accuracy as well as theoretical threshold (vertical line). Treatment **SYM**.

with their signal if signal accuracy is good enough is what underlies the comparatively good information revelation properties highlighted earlier.




(a) RED signal

(b) BLUE signal

Figure 5: The figure shows the vote distribution (red, abstain, blue) conditional on signal accuracy as well as theoretical thresholds (vertical lines) in equilibrium of type II. Treatment **ASYM**

Figure 5 shows how participants vote depending on their signal in treatment **ASYM**. In line with the symmetric equilibrium prediction, if participants receive a BLUE signal they essentially always vote BLUE (panel (b)). There is some abstention for low signal accuracy (below 0.65) and a few votes for RED. Despite the fact that only a small proportion of those receiving a BLUE signal vote RED, this small amount of noise can have a big impact on information revelation as seen above. The reason is that, many more participants do receive a BLUE signal than a RED signal.



(a) RED signal

(b) BLUE signal

Figure 6: The figure shows the vote distribution (red, abstain, blue) conditional on signal accuracy as well as theoretical thresholds (vertical lines) in equilibrium of type II. Treatment **ASYM-COARSE**.

Conditional on receiving a RED signal the majority of participants do *not* vote RED (between 90% if  $q = 0.5$  and around 60% if  $q = 0.9$ ). Around 40-50% of participants vote against their signal, i.e. vote BLUE when their signal was RED. This share is pretty stable across levels of accuracy  $q$  (see also columns (1) and (3) in Table 7). The share of participants abstaining is around 30% for low signal accuracy and decreasing as signal accuracy increases with participants starting to vote RED. Hence, participants with a RED signal vote according to their signal too much if signal accuracy is bad and too little if signal accuracy is good. Hence, rather than a general tendency towards conservatism or base-rate neglect we find that participants do not react enough to the accuracy of their signal.

Figure 6 shows the analogous graph for treatment **ASYM-COARSE**, where signals are less accurate. Behavior conditional on a BLUE signal is very similar to treatment **ASYM** with most participants voting BLUE. Conditional on receiving a RED signal, participants now are even more likely to vote against the signal with almost 80% voting BLUE if  $q = 0.5$ , i.e. if the signal is uninformative. The percentage shrinks for higher levels of  $q$ , but remains substantial at 60% even if  $q = 0.9$ .

Table 5 shows how odds ratios (exponentiated coefficients) from multinomial logit regression on voting outcomes (categorized as “voting with the signal”, “abstaining” or “voting against the signal”) change with the signal accuracy separately for the case where a RED (columns (1)-(3)) or a BLUE (columns (4)-(6)) signal was received. The baseline category is to vote with the signal.<sup>5</sup>

<sup>5</sup>Tables 8 and 7 in Appendix C show OLS regressions on binary outcomes of “Abstention” and “Voting against the signal”, respectively.

	(1)	(2)	(3)	(4)	(5)	(6)
		<i>RED signal</i>		<i>BLUE signal</i>		
	<b>SYM</b>	<b>ASYM</b>	<b>ASYM</b>	<b>SYM</b>	<b>ASYM</b>	<b>ASYM</b>
			<b>COARSE</b>			<b>COARSE</b>
Vote with signal						
Abstain						
q	0.000*** (0.000)	74.460*** (103.500)	53.360 (192.800)	0.005*** (0.000)	0.001 (0.002)	0.097 (0.189)
Constant	135.100*** (101.100)	0.024*** (0.023)	0.010** (0.023)	123.700*** (104.200)	0.754** (0.660)	0.525 (0.633)
Vote against signal						
q	0.044*** (0.051)	0.887 (1.076)	5.681 (12.700)	0.001*** (0.001)	0.025 (0.057)	0.016 (0.058)
Constant	1.816 (1.507)	0.799 (0.624)	0.133 (0.177)	9.781** (11.270)	0.347 (0.538)	0.703 (0.350)
Observations	385	243	181	335	477	539

Standard Errors Eform in parentheses

\*\*\* p<0.01, \*\* p<0.05, \* p<0.1

Table 5: Odds Ratios (exponentiated coefficients) from multinomial logit regression on voting outcomes categorized as “voting with the signal”, “abstaining” or “voting against the signal”. Note: (\*\*\*) significant at 1% level, (\*\*) at 5% level and (\*) at 10% level.

The table shows that conditional on receiving a BLUE signal (columns (4)-(6)), a one-unit increase in  $q$  leads to relative odds to vote against the signal or to abstain that are substantially lower than what they were before. This decline is observed across all treatments, most dramatic in **SYM** (with odds ratios as low as 0.005\*\*\* and 0.001\*\*\*) and least strong and not statistically significant in **ASYM-COARSE**.

Conditional on receiving a RED signal (columns (1)-(3)) the same pattern can be observed in treatment **SYM**: a one-unit increase in  $q$  leads to a substantial decrease in relative odds to vote against the signal or to abstain (relative to voting with the signal). This is not the case in treatment **ASYM**, though. While participants become less likely to vote against RED as signal accuracy increases (exp coefficient 0.088), they do not vote with the signal more, but instead decide to abstain (exp coefficient 74.46\*\*\*).

To sum up, behaviour in the experiment is largely in line with what we expected from theory. Participants are not hesitant to vote against their signal if its quality is low, but they do not react enough to signal accuracy. More informative priors lead to higher efficiency, as expected, but against our expectations, more informative signals do not.

## 4 Further Theoretical Results

### 4.1 Large Elections

In this section we extend our theoretical results by focusing on elections where the number of voters tends to infinite. Our first result is that in large elections the fraction of voters who vote against their signal converges to zero and, moreover, the difference in behavior between those who receive different signals of the same quality also converges to zero.

**Theorem 2.** *The equilibrium in a large election is either Type 1 with  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq \rightarrow 0$  or Type 2 with  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ .*

The first part of the theorem states that in equilibrium of Type 1 we have  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq \rightarrow 0$ . Since in such equilibrium  $q_B^- \leq q_B^+$ , we have then that  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq \rightarrow 0$ . Therefore, the proportion of voters that vote against their signal converges to zero. Note that it may happen that the number of voters who vote against their signal is bounded away from zero, that is, it could be that  $\lim_{N \rightarrow \infty} (N+1) \int_{\frac{1}{2}}^{q_B^-} f(q) dq > \varepsilon$  for some  $\varepsilon > 0$ . However, the number of voters voting against their signal in the population is insignificant compared with the number of voters who vote according to their signal.

Another implication of first part of the theorem is that the difference in behavior between those who receive signal  $R$  or  $B$  converges to zero. A voter who receives signal  $R$  always votes for  $R$  while the fraction of voters who do not vote for  $B$  when they receive signal  $B$  is  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq$ , which converges to zero.

The second part of the theorem states that in equilibrium of Type 2 we have  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ . Again this implies that the difference in behavior between those who receive signal  $R$  or  $B$  converges to zero. Therefore, Theorem 2 implies that when the number of voters tends to infinity the fraction of voters who vote against their signal ( $\int_{\frac{1}{2}}^{q_B^-} f(q) dq$ ) and the fraction of voters whose behavior depend on the specific signal received ( $\int_{q_R^+}^{q_B^+} f(q) dq$ ) vanishes in the limit. That is, as the number of voters increases the effect of an asymmetric common prior ( $p > \frac{1}{2}$ ) vanishes in the limit and the results in McMurray (2013) apply. (i.e. the equilibrium is characterized by a cut-point  $q$  that determines who abstains and who votes for his signal independently on the particular signal received).

The reason behind the result in Theorem 2 is the following. Assume that  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq$  does not converge to zero. In this case if a voter is pivotal then it must be that in proportion more voters received signal  $B$  than  $R$ : as  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq$  does not converge to zero, not all voters who receive signal  $B$  vote for  $B$  yet all voters who receive signal  $R$  vote for  $R$ . If more voters

receive signal  $B$  than  $R$  then since the average signal quality  $\mu$  is greater than  $\frac{1}{2}$  by law of large numbers the state of nature is  $B$  with probability one. This implies that all voters should vote for  $B$ , contradicting the fact that all voters who receive signal  $R$  prefer to vote for  $R$ .

A similar argument shows that as the number of voters grows large in equilibrium of Type 2 we must have  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ . If  $\int_{q_R^+}^{q_B^+} f(q) dq$  does not converge to zero then if a voter is pivotal it must be that a greater proportion of voters receive signal  $B$  than signal  $R$ . This is because a higher fraction of those voters who receive signal  $B$  compared to those who receive signal  $R$  abstain. Law of large number then means that the state is  $B$  with probability one, which implies that all voters should vote for  $B$ . This represents a contradiction to the characterization of equilibrium of Type 2.

In this paper we consider only symmetric strategies. The fact that an equilibrium where some voters vote against their signal exists in symmetric strategies implies that it also exists when non-symmetric strategies are considered. Moreover, the fact that we consider only symmetric strategies is without loss of generality in case there is a large number of voters. This is because in large elections the probability that the vote of a specific voter determines the outcome is zero. Thus, all voters face the same distribution of strategies played by other voters. Given that voters are never (with probability zero) indifferent between the available options (voting for either candidate and abstaining), all voters chose the same strategy conditional on the signal received and its quality. That is, voters follow symmetric strategies. For more on this see McMurray (2013).

Our final result states that in large elections the best candidate wins with probability one. This result is in line with the Condorcet Jury Theorem and the findings in previous literature (see for instance Feddersen and Pesendorfer (1996, 1997, 1999) and McMurray (2013)).

**Proposition 1.** *The equilibrium in a large election is such that the best candidate wins with probability one.*

Given the result in Theorem 2, whether a voter chooses to vote or to abstain depends on the quality of his signal, not on the value of the signal itself. Thus, for a given state of nature and given level of abstention, the best candidate is expected to receive a share  $\mu$  of the votes while the other candidate is expected to receive a share  $1 - \mu$  of the votes. Since  $\mu > \frac{1}{2}$  law of large numbers implies that the best candidate wins with probability one.



## 5 Conclusions

We presented a common value election setting where voters have private information of different qualities. We showed both theoretically and experimentally that voters may have incentives to vote against their private information, even if such private information is useful, all have the same preferences, and abstention is allowed. Moreover, we found that elections do not generally aggregate information efficiently. On top of that, experimental subjects used their private information more than what is rational. This produced the unexpected result that lower quality of information is better; the reduction in the quality of private information made subjects trust their information less, which reduced the efficiency loss due to subjects following their private information too often.

We also found that the behavior of voters when they receive information of different qualities resembles the behavior of voters who receive information of the same quality but that have different preferences (i.e. they may be biased towards either candidate). This suggests that the observed different biases towards candidates need not be the result of different preferences and it could be the result of different rational beliefs about who is the best candidate.

## References

- Austen-Smith, D. and J. S. Banks (1996). Information aggregation, rationality, and the Condorcet jury theorem. *The American Political Science Review* 90(1), 34–45.
- Battaglini, M., R. Morton, and T. Palfrey (2008). Information aggregation and strategic abstention in large laboratory elections. *American Economic Review* 98(2), 194–200.
- Battaglini, M., R. Morton, and T. Palfrey (2010). The swing voter’s curse in the laboratory. *Review of Economic Studies* 77, 61–89.
- Ben-Yashar, R. and I. Milchtaich (2007). First and second best voting rules in committees. *Social Choice and Welfare* 29, 453–486.
- Condorcet, M. d. (1785). Essai sur la application del analyse à la probabilité des décisions rendues à la probabilité des voix. *De l’Impremiere Royale, Paris*.
- Coughlan, P. (2000). In defense of unanimous jury verdicts: mistrials, communication and strategic voting. *American Political Science Review* 94, 375–393.
- Erev, I., D. Shimonowitch, A. Schurr, and R. Hertwig (2008). Base rates: how to make the

- intuitive mind appreciate or neglect them. In H. Plessner, C. Betsch, and T. Betsch (Eds.), *Intuition in Judgement and Decision-Making*, pp. 135–148. NJ: Lawrence Erlbaum.
- Feddersen, T. and W. Pesendorfer (1996). The swing voters curse. *The American Economic Review* 86(3), 408–424.
- Feddersen, T. and W. Pesendorfer (1997). Voting behavior and information aggregation in elections with private information. *Econometrica* 65(5), 1029–1058.
- Feddersen, T. and W. Pesendorfer (1998). Abstention in elections with asymmetric information and diverse preferences. *The American Political Science Review* 93(2), 381–398.
- Fischbacher, U. (2007). z-tree: Zurich toolbox for ready-made economic experiments. *Experimental Economics* 10(2), 171–178.
- Grether, D. (1980). Bayes rule as a descriptive model: the representativeness heuristic. *Quarterly Journal of Economics* 95, 537–557.
- Guarnaschelli, S., R. McKelvey, and T. Palfrey (2000). An experimental study of jury decision rules. *American Political Science Review* 94(2), 407–423.
- Kahnemann, D. and A. Tversky (1972). Subjective probability: a judgement of representativeness. *Cognitive Psychology* 3, 430–454.
- Krishna, V. and J. Morgan (2012). Voluntary voting: Costs and benefits. *Journal of Economic Theory* 147, 2083–2123.
- McMurray, J. (2013). Aggregating information by voting: The wisdom of experts versus the wisdom of the masses. *Review of Economic Studies* 80(1), 277–312.
- McMurray, J. C. (2010). Empirical evidence of strategic voter abstention. mimeo.
- Morton, R. and J. Tyran (2011). Let the experts decide? asymmetric information, abstention and coordination in standing committees. *Games and Economic Behavior* 72, 485–509.
- Myatt, D. (2012). A rational choice theory of voter turnout. *working paper*.
- Myerson, R. B. (1998). Extended poisson games and the condorcet jury theorem. *Games and Economic Behavior* 25, 111–131.
- Myerson, R. B. (2000). Large poisson games. *Journal of Economic Theory* 94, 7–45.
- Nunez, M. (2010). Condorcet consistency of approval voting: A counter example in large poisson games. *Journal of Theoretical Politics* 22, 64–84.

- Oliveros, S. (2013). Abstention, ideology and information acquisition. *Journal of Economic Theory* 148, 871–902.
- Rivas, J. and C. Rodríguez-Álvarez (2012). Deliberation, leadership and information aggregation. *University of Leicester Working Paper 12/16*.
- Ward, E. (1982). Conservatism in human information processing. In D. Kahnemann, P. Slovic, and A. Tversky (Eds.), *Judgment under uncertainty: Heuristics and biases*. New York: Cambridge University Press.

## A Appendix: Proofs

The following lemma is used in the proof of Theorem 1.

**Lemma 1.** *The best response of any voter  $i$  against any strategy  $v$  played by the other  $N$  voters is given  $v'$ , which is characterized by four cutpoints  $q_R^-, q_R^+, q_B^-$  and  $q_B^+$  in  $[\frac{1}{2}, 1]$  such that*

$$v'(\theta_i, q_i) = \begin{cases} R & \text{if either } \sigma_i = R \text{ and } q_i \geq q_R^+ \text{ or } \sigma_i = B \text{ and } q_i < q_B^-, \\ B & \text{if either } \sigma_i = R \text{ and } q_i < q_R^- \text{ or } \sigma_i = B \text{ and } q_i \geq q_B^+, \\ \emptyset & \text{otherwise.} \end{cases}$$

*Proof.* Take any arbitrary voter  $i$  and assume all voters except  $i$  use strategy  $v$ . Consider equations (1), (2) and (3) and assume that  $\sigma_i = R$ . We have that both  $Eu(R, v) - Eu(B, v)$  and  $Eu(R, v) - Eu(\emptyset, v)$  are increasing in  $q_i$ . Therefore, there exists a  $x \in [0, 1]$  such that both equations are positive and voter  $i$  votes for  $R$  whenever  $q_i \geq x$ . Since  $q_i \in [\frac{1}{2}, 1]$  if we define  $q_R^+ = \max\{\frac{1}{2}, x\}$  we have that voter  $i$  votes for  $R$  whenever  $q_i \geq q_R^+$ .

Moreover, both  $Eu(B, v) - Eu(R, v)$  and  $Eu(B, v) - Eu(\emptyset, v)$  are decreasing in  $q_i$ . Therefore, there exists a  $y$  with  $0 \leq y \leq x$  such that both equations are positive and voter  $i$  votes for  $B$  whenever  $q_i < y$ . If we define  $q_R^- = \max\{\frac{1}{2}, y\}$  we have that voter  $i$  votes for  $R$  whenever  $q_i < q_R^-$ .

The final possibility is that both  $Eu(R, v) - Eu(\emptyset, v)$  and  $Eu(B, v) - Eu(\emptyset, v)$  are negative, which can happen if and only if  $q_i \in [y, x)$  or, in other words,  $q_i \in [q_R^-, q_R^+)$ . In this case, voter  $i$  prefers to abstain.

A similar reasoning when  $\sigma_i = B$  leads to the conclusion in the lemma.  $\square$

### *Proof of Theorem 1. An Equilibrium Exists*

First we demonstrate existence. Given the result in Lemma 1, we know that for any strategy  $v$  employed by the other  $N$  voters every voter employs a strategy that is characterized by four cutpoints  $q_R^-, q_R^+, q_B^-$  and  $q_B^+$ . Define the function  $\phi : [\frac{1}{2}, 1]^4 \rightarrow [\frac{1}{2}, 1]^4$  where  $\phi(q_R^-, q_R^+, q_B^-, q_B^+)$  is the best response of any voter to a situation where all other  $N$  voters employ an strategy characterized the four cutpoints  $q_R^-, q_R^+, q_B^-, q_B^+$ . We have to prove that  $\phi$  has a fixed point. By the fixed point theorem, since the set  $[\frac{1}{2}, 1]^4$  is convex and compact in the Euclidean space we are left to show that  $\phi$  is continuous.

When  $N$  voters are using strategy  $v$  characterized by the four cutpoints  $q_R^-, q_R^+, q_B^-$  and

$q_B^+$  we have that

$$\begin{aligned}
\pi_t(v, R) &= \sum_{s_R(R)=0}^{\lfloor \frac{N}{2} \rfloor} \sum_{s_B(R)=0}^{\lfloor \frac{N}{2} \rfloor - s_R(R)} \sum_{s_R(\emptyset)=0}^{N - 2(s_R(R) + s_B(R))} \sum_{s_R(B)=0}^{s_R(R) + s_B(R)} \\
&\quad \frac{N!}{s_R(R)! s_B(R)! s_R(\emptyset)! s_R(B)! (s_R(R) + s_B(R) - s_R(B))! (N - 2(s_R(R) + s_B(R)) - s_R(\emptyset))!} \\
&\quad \times \left[ \int_{q_R^+}^1 qf(q) dq \right]^{s_R(R)} \left[ \int_{\frac{1}{2}}^{q_B^-} (1-q)f(q) dq \right]^{s_B(R)} \\
&\quad \times \left[ \int_{q_R^-}^{q_R^+} qf(q) dq \right]^{s_R(\emptyset)} \left[ \int_{\frac{1}{2}}^{q_R^-} qf(q) dq \right]^{s_R(B)} \\
&\quad \times \left[ \int_{q_B^+}^1 (1-q)f(q) dq \right]^{s_R(R) + s_B(R) - s_R(B)} \left[ \int_{q_B^-}^{q_B^+} (1-q)f(q) dq \right]^{N - 2(s_R(R) + s_B(R)) - s_R(\emptyset)} \quad (4)
\end{aligned}$$

Since  $F(q) = \int_{\frac{1}{2}}^q f(q) dq$  we have that  $F$  is continuous and, because it is a cumulative density function, it is bounded in  $[0, 1]$ . Therefore,  $F$  is integrable and moreover continuous with respect to the integration limits. Thus,  $\int qf(q) dq = qF(q) - \int F(q) dq$  is continuous with respect to the integration limits. As a result,  $\pi_t(v, R)$  is continuous with respect to the cutpoints  $q_R^-, q_R^+, q_B^-$  and  $q_B^+$ .

It can be shown in a similar fashion that  $\pi_t(v, s), \pi_B(v, s)$  and  $\pi_R(v, s)$  are continuous with respect to the cutpoints  $q_R^-, q_R^+, q_B^-$  and  $q_B^+$  for all  $s \in \{R, B\}$ . Hence, we have that  $Eu(R, v) - Eu(B, v), Eu(R, v) - Eu(\emptyset, v)$  and  $Eu(R, v) - Eu(\emptyset, v)$  are continuous with respect to the cutpoints  $q_R^-, q_R^+, q_B^-$  and  $q_B^+$ . Thus,  $\phi$  is continuous as we wanted to show.

### Equilibrium is of Two Types

Given the result in Lemma 1, any equilibrium is characterized by the four threshold values  $q_R^-, q_R^+, q_B^-$  and  $q_B^+$ . Assume that  $q_B^- > \frac{1}{2}$ , then we have that  $Eu(R, v) - Eu(B, v) > 0$  and  $Eu(R, v) - Eu(\emptyset, v) > 0$  for all  $i$  with  $\sigma_i = B$  and  $q_i \in [\frac{1}{2}, q_B^-)$ , which implies that  $Eu(R, v) - Eu(B, v) > 0$  and  $Eu(R, v) - Eu(\emptyset, v) > 0$  for all  $i$  with  $\sigma_i = R$  and  $q_i \in [\frac{1}{2}, q_B^-)$ . This means that  $q_R^-, q_R^+ = \frac{1}{2}$ , which leads to equilibrium of Type 1 in the proposition.

Assume now that  $q_B^- = \frac{1}{2}$  and  $q_B^+ > \frac{1}{2}$ . In this case we have that  $Eu(B, v) - Eu(\emptyset, v) < 0$  for all  $i$  with  $\sigma_i = B$  and  $q_i \in [\frac{1}{2}, q_B^+)$ , which implies that  $Eu(B, v) - Eu(\emptyset, v) < 0$  for all  $i$  with  $\sigma_i = R$  and  $q_i \in [\frac{1}{2}, q_B^+)$ . This means that  $q_R^- = \frac{1}{2}$ , which leads to equilibrium of type 2 in the proposition.

Finally, assume that  $q_B^- = q_B^+ = \frac{1}{2}$ . We proceed by showing that  $\pi_t(v, R) + \pi_B(v, R) \geq \pi_t(v, B) + \pi_B(v, B)$ . If this were true, and since  $q_B^+ = \frac{1}{2}$  implies that  $Eu(B, v) - Eu(\emptyset, v) \geq 0$  for all  $i$  with  $\sigma_i = B$  and  $q_i \in [\frac{1}{2}, 1]$ , equation (3) together with the fact that  $p \geq \frac{1}{2}$  implies

that  $\pi_t(v, B) + \pi_B(v, B) \geq \pi_t(v, R) + \pi_B(v, R)$ , which would represent a contradiction (unless  $q_B^- = q_B^+ = q_R^- = q_R^+ = p = \frac{1}{2}$ , which is an equilibrium of either Type in the proposition).

First we show that  $\pi_t(v, R) - \pi_t(v, B) \geq 0$  for all  $\frac{1}{2} \leq q_R^- \leq q_R^+ \leq 1$ . Note that

$$\begin{aligned}\pi_t(v, R) &= \sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \sum_{r=0}^j \frac{N!}{j!(j-r)!r!(N-2j)!} \\ &\quad \left[ \int_{q_R^+}^1 qf(q) dq \right]^j \left[ \int_{\frac{1}{2}}^{q_R^-} qf(q) dq \right]^{j-r} \left[ \int_{\frac{1}{2}}^1 (1-q)f(q) dq \right]^r \left[ \int_{q_R^-}^{q_R^+} qf(q) dq \right]^{N-2j}, \\ \pi_t(v, B) &= \sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \sum_{r=0}^j \frac{N!}{j!(j-r)!r!(N-2j)!} \\ &\quad \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right]^j \left[ \int_{\frac{1}{2}}^{q_R^-} (1-q)f(q) dq \right]^{j-r} \left[ \int_{\frac{1}{2}}^1 qf(q) dq \right]^r \left[ \int_{q_R^-}^{q_R^+} (1-q)f(q) dq \right]^{N-2j}.\end{aligned}$$

Given that  $q_i \geq \frac{1}{2}$  for all voter  $i$  we have that

$$\begin{aligned}\pi_t(v, R) - \pi_t(v, B) &\geq \sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \sum_{r=0}^j \frac{N!}{j!(j-r)!r!(N-2j)!} \\ &\quad \left( \left[ \int_{q_R^+}^1 qf(q) dq \right]^j \left[ \int_{\frac{1}{2}}^1 (1-q)f(q) dq \right]^r \right. \\ &\quad \left. - \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right]^j \left[ \int_{\frac{1}{2}}^1 qf(q) dq \right]^r \right).\end{aligned}$$

Thus, if  $q_R^+ = \frac{1}{2}$  or  $q_R^+ = 1$  then  $\pi_t(v, R) - \pi_t(v, B) \geq 0$ . Consider now the cases where  $q_R^+ \in (\frac{1}{2}, 1)$ . Using once more that  $q_i \geq \frac{1}{2}$  for all  $i$ , a necessary condition for  $\pi_t(v, R) - \pi_t(v, B) \geq 0$  is that

$$\left[ \int_{q_R^+}^1 qf(q) dq \right]^r \left[ \int_{\frac{1}{2}}^1 (1-q)f(q) dq \right]^r \geq \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right]^r \left[ \int_{\frac{1}{2}}^1 qf(q) dq \right]^r.$$

This can be written as

$$\begin{aligned}\left[ \int_{q_R^+}^1 qf(q) dq \right] \left[ \int_{q_R^+}^1 (1-q)f(q) dq + \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right] &\geq \\ \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right] \left[ \int_{q_R^+}^1 qf(q) dq + \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]. &\end{aligned}$$

In other words,

$$\begin{aligned}
\frac{\int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq}{\int_{q_R^+}^1 (1-q)f(q) dq} &\geq \frac{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq}{\int_{q_R^+}^1 qf(q) dq}, \\
\frac{\int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq}{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq} &\geq \frac{\int_{q_R^+}^1 (1-q)f(q) dq}{\int_{q_R^+}^1 qf(q) dq}, \\
\frac{\int_{\frac{1}{2}}^{q_R^+} f(q) dq}{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq} &\geq \frac{\int_{q_R^+}^1 f(q) dq}{\int_{q_R^+}^1 qf(q) dq}, \\
\frac{\int_{q_R^+}^1 qf(q) dq}{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq} &\geq \frac{\int_{q_R^+}^1 f(q) dq}{\int_{\frac{1}{2}}^{q_R^+} f(q) dq}.
\end{aligned}$$

Since it is true that

$$\begin{aligned}
\frac{\int_{q_R^+}^1 qf(q) dq}{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq} &\geq \frac{\int_{q_R^+}^1 q_R^+ f(q) dq}{\int_{\frac{1}{2}}^{q_R^+} q_R^+ f(q) dq} \\
&= \frac{\int_{q_R^+}^1 f(q) dq}{\int_{\frac{1}{2}}^{q_R^+} f(q) dq},
\end{aligned}$$

we have that  $\pi_t(v, R) - \pi_t(v, B) \geq 0$ .

Proceeding in a similar fashion, it can be shown that  $\pi_B(v, R) - \pi_B(v, B) \geq 0$ . Thus, we have that  $\pi_t(v, R) - \pi_B(v, R) \geq \pi_t(v, B) - \pi_B(v, B)$  as required.

### **In Equilibrium of Type 1** $q_B^- \leq p$

We can use the algebra from the previous part of the proof to show that in equilibrium of Type 1  $\pi_t(v, R) - \pi_t(v, B) \leq 0$  and  $\pi_R(v, B) - \pi_R(v, R) \geq 0$  for all  $\frac{1}{2} \leq q_B^- \leq q_B^+ \leq 1$ . Hence, equation (2) together with the definition of  $q_B^-$  implies  $p(1 - q_B^-) \geq (1 - p)q_B^-$ , which in turn implies  $q_B^- \leq p$ .

### **In Equilibrium of Type 2** $q_B^+ \geq q_R^+$

Next we prove that in any equilibrium of type 2 it must be that  $q_B^+ \geq q_R^+$ . Assume the

opposite,  $q_R^+ > q_B^+$ . Note that in any Type 2 equilibrium we have that

$$\begin{aligned} \pi_t(v, R) &= \sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \left[ \int_{q_R^+}^1 qf(q) dq \right]^j \left[ \int_{q_B^+}^1 (1-q)f(q) dq \right]^j \\ &\quad \times \sum_{r=0}^{N-2j} \frac{N!}{j!r!(N-2j-r)!} \\ &\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^r \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^{N-2j-r}. \end{aligned}$$

Thus, it is true that

$$\begin{aligned} \pi_t(v, R) &= \sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \left[ \int_{q_R^+}^1 qf(q) dq \right]^j \left[ \int_{q_B^+}^1 (1-q)f(q) dq \right]^j \\ &\quad \times \sum_{k=0}^{\lfloor \frac{N-2j}{2} \rfloor} \frac{N!}{j!k!(N-2j-k)!} \\ &\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^k \\ &\quad \times \left( \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^{N-2j-2k} + \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^{N-2j-2k} \right), \end{aligned} \quad (5)$$

$$\begin{aligned} \pi_t(v, B) &= \sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right]^j \left[ \int_{q_B^+}^1 qf(q) dq \right]^j \\ &\quad \times \sum_{k=0}^{\lfloor \frac{N-2j}{2} \rfloor} \frac{N!}{j!k!(N-2j-k)!} \\ &\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^k \\ &\quad \times \left( \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^{N-2j-2k} + \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^{N-2j-2k} \right). \end{aligned} \quad (6)$$

We now show that  $\pi_t(v, R) - \pi_t(v, B) \geq 0$  in three steps. First, we have that  $q_R^+ > q_B^+$  implies

$$\left[ \int_{q_R^+}^1 qf(q) dq \right]^j \left[ \int_{q_B^+}^1 (1-q)f(q) dq \right]^j \geq \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right]^j \left[ \int_{q_B^+}^1 qf(q) dq \right]^j \quad (7)$$



for all  $j \in \{0, 1, \dots\}$  if and only if

$$\int_{q_R^+}^1 qf(q)dq \left[ \int_{q_R^+}^1 (1-q)f(q)dq + \int_{q_B^+}^{q_R^+} (1-q)f(q)dq \right] \geq \int_{q_R^+}^1 (1-q)f(q)dq \left[ \int_{q_R^+}^1 qf(q)dq + \int_{q_B^+}^{q_R^+} qf(q)dq \right].$$

which can be rewritten as

$$\begin{aligned} \int_{q_R^+}^1 qf(q)dq \int_{q_B^+}^{q_R^+} (1-q)f(q)dq &\geq \int_{q_R^+}^1 (1-q)f(q)dq \int_{q_B^+}^{q_R^+} qf(q)dq, \\ \frac{\int_{q_B^+}^{q_R^+} (1-q)f(q)dq}{\int_{q_B^+}^{q_R^+} qf(q)dq} &\geq \frac{\int_{q_R^+}^1 (1-q)f(q)dq}{\int_{q_R^+}^1 qf(q)dq}, \\ \frac{\int_{q_B^+}^{q_R^+} f(q)dq}{\int_{q_B^+}^{q_R^+} qf(q)dq} &\geq \frac{\int_{q_R^+}^1 f(q)dq}{\int_{q_R^+}^1 qf(q)dq}, \\ \frac{\int_{q_R^+}^1 qf(q)dq}{\int_{q_B^+}^{q_R^+} qf(q)dq} &\geq \frac{\int_{q_R^+}^1 f(q)dq}{\int_{q_B^+}^{q_R^+} f(q)dq}, \end{aligned}$$

which given that

$$\begin{aligned} \frac{\int_{q_R^+}^1 qf(q)dq}{\int_{q_B^+}^{q_R^+} qf(q)dq} &\geq \frac{\int_{q_R^+}^1 q_R^+ f(q)dq}{\int_{q_B^+}^{q_R^+} q_R^+ f(q)dq}, \\ &= \frac{\int_{q_R^+}^1 f(q)dq}{\int_{q_B^+}^{q_R^+} f(q)dq}, \end{aligned}$$

proves that equation (7) holds true when  $q_R^+ > q_B^+$ .

Second, we have that  $q_R^+ > q_B^+$  implies

$$\left[ \int_{\frac{1}{2}}^{q_R^+} qf(q)dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q)dq \right]^k \geq \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q)dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q)dq \right]^k \quad (8)$$

for all  $k \in \{0, 1, \dots\}$  if and only if

$$\begin{aligned} \frac{\int_{\frac{1}{2}}^{q_B^+} f(q) dq}{\int_{\frac{1}{2}}^{q_R^+} f(q) dq} &\geq \frac{\int_{\frac{1}{2}}^{q_B^+} qf(q) dq}{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq}, \\ \frac{\int_{\frac{1}{2}}^{q_R^+} qf(q) dq}{\int_{\frac{1}{2}}^{q_B^+} qf(q) dq} &\geq \frac{\int_{\frac{1}{2}}^{q_R^+} f(q) dq}{\int_{\frac{1}{2}}^{q_B^+} f(q) dq}, \\ \frac{\int_{q_B^+}^{q_R^+} qf(q) dq}{\int_{\frac{1}{2}}^{q_B^+} qf(q) dq} &\geq \frac{\int_{q_B^+}^{q_R^+} f(q) dq}{\int_{\frac{1}{2}}^{q_B^+} f(q) dq}, \end{aligned}$$

which given that

$$\begin{aligned} \frac{\int_{q_B^+}^{q_R^+} qf(q) dq}{\int_{\frac{1}{2}}^{q_B^+} qf(q) dq} &\geq \frac{\int_{q_B^+}^{q_R^+} q^+ f(q) dq}{\int_{\frac{1}{2}}^{q_B^+} q^+ f(q) dq}, \\ &= \frac{\int_{q_B^+}^{q_R^+} f(q) dq}{\int_{\frac{1}{2}}^{q_B^+} f(q) dq}, \end{aligned}$$

proves that equation (8) holds true when  $q_R^+ > q_B^+$ .

Third, we have that  $q_R^+ > q_B^+$  implies

$$\begin{aligned} \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^m + \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^m &\geq \\ \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^m + \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^m &\end{aligned} \quad (9)$$

for all  $m \in \{0, 1, \dots\}$  if and only if

$$\begin{aligned} \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq + \int_{q_B^+}^{q_R^+} qf(q) dq \right]^m - \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^m &\geq \\ \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq + \int_{q_B^+}^{q_R^+} (1-q)f(q) dq \right]^m - \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^m &\end{aligned}$$

which is always true for  $m = 0$  and true for  $m \in \{1, 2, \dots\}$  if and only if

$$\begin{aligned} \sum_{l=1}^m \binom{m}{l} \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^{m-l} \left[ \int_{q_B^+}^{q_R^+} qf(q) dq \right]^l &\geq \\ \sum_{l=1}^m \binom{m}{l} \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^{m-l} \left[ \int_{q_B^+}^{q_R^+} (1-q)f(q) dq \right]^l &\end{aligned}$$

Since the expression above is true we have that  $q_R^+ > q_B^+$  implies equation (9) as required.

Therefore, we have shown that  $q_R^+ > q_B^+$  implies equations (7), (8) and (9) are true. Hence, from equations (5) and (6) we have that  $q_R^+ > q_B^+$  implies  $\pi_t(v, R) - \pi_t(v, B) \geq 0$ .

Equations (2) and (3) together with the fact that  $q_R^+ > q_B^+$  and  $p \geq \frac{1}{2}$  imply that

$$\begin{aligned} q_R^+ (\pi_t(v, R) + \pi_R(v, R)) &\leq (1 - q_R^+) (\pi_t(v, B) + \pi_R(v, B)), \\ q_R^+ (\pi_t(v, B) + \pi_B(v, B)) &> (1 - q_R^+) (\pi_t(v, R) + \pi_B(v, R)). \end{aligned}$$

Given that, as we have just shown,  $q_R^+ > q_B^+$  implies  $\pi_t(v, R) - \pi_t(v, B) \geq 0$ , the two expressions above imply

$$(1 - q_R^+) (\pi_R(v, B) - \pi_B(v, R)) > q_R^+ (\pi_R(v, R) - \pi_B(v, B)). \quad (10)$$

Note now that

$$\begin{aligned} \pi_B(v, R) &= \sum_{j=1}^{\lfloor \frac{N}{2} \rfloor} \left[ \int_{q_R^+}^1 qf(q) dq \right]^j \left[ \int_{q_B^+}^1 (1-q)f(q) dq \right]^{j-1} \\ &\quad \times \sum_{k=0}^{\lfloor \frac{N-2j+1}{2} \rfloor} \frac{N!}{j!j!k!(N-2j+1-k)!} \\ &\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^k \\ &\quad \times \left( \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^{N-2j+1-2k} + \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^{N-2j+1-2k} \right), \\ \pi_R(v, B) &= \sum_{j=1}^{\lfloor \frac{N}{2} \rfloor} \left[ \int_{q_R^+}^1 (1-q)f(q) dq \right]^{j-1} \left[ \int_{q_B^+}^1 qf(q) dq \right]^j \\ &\quad \times \sum_{k=0}^{\lfloor \frac{N-2j+1}{2} \rfloor} \frac{N!}{j!j!k!(N-2j+1-k)!} \\ &\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^k \\ &\quad \times \left( \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^{N-2j+1-2k} + \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^{N-2j+1-2k} \right), \end{aligned}$$

and similarly for  $\pi_R(v, R)$  and  $\pi_B(v, B)$ . Define

$$\begin{aligned}
K_R &= \sum_{k=0}^{\lfloor \frac{N-2j+1}{2} \rfloor} \frac{N!}{j!j!k!(N-2j-k+1)!} \\
&\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^k \\
&\quad \times \left( \left[ \int_{\frac{1}{2}}^{q_R^+} qf(q) dq \right]^{N-2j+1-2k} + \left[ \int_{\frac{1}{2}}^{q_B^+} (1-q)f(q) dq \right]^{N-2j+1-2k} \right), \\
K_B &= \sum_{k=0}^{\lfloor \frac{N-2j+1}{2} \rfloor} \frac{N!}{j!j!k!(N-2j-k+1)!} \\
&\quad \times \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^k \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^k \\
&\quad \times \left( \left[ \int_{\frac{1}{2}}^{q_R^+} (1-q)f(q) dq \right]^{N-2j+1-2k} + \left[ \int_{\frac{1}{2}}^{q_B^+} qf(q) dq \right]^{N-2j+1-2k} \right).
\end{aligned}$$

Then, equations (8) and (9) imply  $K_R \geq K_B$ . Moreover, as  $q_R^+ > q_B^+$  implies equation (7), we have that

$$\begin{aligned}
(1 - q_R^+) (\pi_R(v, B) - \pi_B(v, R)) &\leq (1 - q_R^+) \left( \int_{q_B^+}^1 qf(q) dq - \int_{q_R^+}^1 qf(q) dq \right) K_B, \\
q_R^+ (\pi_R(v, R) - \pi_B(v, B)) &\geq q_R^+ \left( \int_{q_B^+}^1 (1-q)f(q) dq - \int_{q_R^+}^1 (1-q)f(q) dq \right) K_R.
\end{aligned}$$

This means that equation (10) holds only if

$$\begin{aligned}
(1 - q_R^+) \left( \int_{q_B^+}^1 qf(q) dq - \int_{q_R^+}^1 qf(q) dq \right) K_B &\geq q_R^+ \left( \int_{q_B^+}^1 (1-q)f(q) dq - \int_{q_R^+}^1 (1-q)f(q) dq \right) K_R, \\
(1 - q_R^+) \left( \int_{q_B^+}^{q_R^+} qf(q) dq \right) &\geq q_R^+ \left( \int_{q_B^+}^{q_R^+} (1-q)f(q) dq \right), \\
\left( \int_{q_B^+}^{q_R^+} qf(q) dq \right) &\geq \left( \int_{q_B^+}^{q_R^+} q_R^+ f(q) dq \right),
\end{aligned}$$

holds. However, given that  $q_R^+ > q_B^+$  the expression above is false. This leads to a contradiction, which means that the claim  $q_R^+ > q_B^+$  is false as required.  $\square$

The following Lemma from Feddersen and Pessendorfer (1996) is used in the proof of the Theorem 2.

**Lemma 2** (Lemma 0 in Feddersen and Pessendorfer (1996)). *Let  $(a_N, b_N, c_N)_{N=1}^{\infty}$  a sequence that satisfies  $(a_N, b_N, c_N) \in [0, 1]^3$  and  $a_N < b_N - \delta$  and  $\delta < c_N$  for all  $N$  and some  $\delta > 0$ . Then, for  $i = 0, 1$  as  $N \rightarrow \infty$*

$$\frac{\sum_{j=0}^{\frac{N}{2}-i} \frac{N!}{(j+i)!j!(N-2j-i)!} c_N^{N-2j-i} a_N^j}{\sum_{j=0}^{\frac{N}{2}-i} \frac{N!}{(j+i)!j!(N-2j-i)!} c_N^{N-2j-i} b_N^j} \rightarrow 0.$$

*Proof of Theorem 2. Equilibrium of Type 1*

First we show that as  $N \rightarrow \infty$  the equilibrium of Type 1 is such that  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq \rightarrow 0$ . Assume for now that there exists a  $\rho > 0$  such that  $\int_{q_B^+}^1 f(q) dq > \rho$  for all  $N$  and consider an equilibrium of Type 1 and assume that there exists a  $\varepsilon > 0$  such that either  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq \geq \varepsilon$  or  $\int_{q_B^-}^{q_B^+} f(q) dq \geq \varepsilon$  for all  $N$ . We have that there exists a  $\delta_1 > 0$  such that  $\sigma_{QQ}\sigma_{AQ} - \delta_1 > \sigma_{AA}\sigma_{QA}$  if and only if

$$\begin{aligned} & \int_{q_B^+}^1 qf(q) dq \left( \int_{\frac{1}{2}}^1 (1-q)f(q) dq + \int_{\frac{1}{2}}^{q_B^-} qf(q) dq \right) - \delta_1 > \\ & \int_{q_B^+}^1 (1-q)f(q) dq \left( \int_{\frac{1}{2}}^1 qf(q) dq + \int_{\frac{1}{2}}^{q_B^-} (1-q)f(q) dq \right), \\ & \int_{q_B^+}^1 qf(q) dq \left( \int_{q_B^-}^1 (1-q)f(q) dq + \int_{\frac{1}{2}}^{q_B^-} f(q) dq \right) - \delta_1 > \\ & \int_{q_B^+}^1 (1-q)f(q) dq \left( \int_{q_B^-}^1 qf(q) dq + \int_{\frac{1}{2}}^{q_B^-} f(q) dq \right), \\ & \int_{q_B^+}^1 qf(q) dq \left( \int_{q_B^-}^{q_B^+} (1-q)f(q) dq + \int_{\frac{1}{2}}^{q_B^-} f(q) dq \right) - \delta_1 > \\ & \int_{q_B^+}^1 (1-q)f(q) dq \left( \int_{q_B^-}^{q_B^+} qf(q) dq + \int_{\frac{1}{2}}^{q_B^-} f(q) dq \right). \end{aligned}$$

R necessary condition for this is

$$\begin{aligned} & \int_{q_B^+}^1 (q - q_B^+) f(q) dq \int_{q_B^-}^{q_B^+} f(q) dq + \int_{q_B^+}^1 (2q - 1) f(q) dq \int_{\frac{1}{2}}^{q_B^-} f(q) dq - \delta_1 > 0, \\ & \int_{q_B^+}^1 (q - q_B^+) f(q) dq \int_{q_B^-}^{q_B^+} f(q) dq + (2q_B^- - 1) \int_{q_B^+}^1 f(q) dq \int_{\frac{1}{2}}^{q_B^-} f(q) dq - \delta_1 > 0. \end{aligned}$$

By assumption  $\int_{q_B^+}^1 f(q) dq > \rho$  for some  $\rho > 0$ . Therefore, if  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq \geq \varepsilon$  then  $q_B^- \geq F^{-1}(\varepsilon)$  and the expression above is true for any  $\delta_1 \in (0, (2F^{-1}(\varepsilon) - 1)\rho\varepsilon)$ .<sup>6</sup>

<sup>6</sup> $F^{-1}$  exists because  $f$  is integrable and, hence,  $F$  is continuous.

Assume  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq < \varepsilon$ , which implies that  $\int_{q_B^-}^{q_B^+} f(q) dq \geq \varepsilon$ . Note that  $\int_{q_B^+}^1 f(q) dq > \rho$  for some  $\rho > 0$  implies that for all  $\bar{\rho} \in (0, \rho)$  there exists a  $\beta > 0$  such that  $\int_{q_B^+ + \beta}^1 f(q) dq > \bar{\rho}$ , fix such  $\rho$  and consider its corresponding  $\beta$ . Thus, a necessary condition for  $\sigma_{QQ}\sigma_{AQ} - \delta_1 > \sigma_{AA}\sigma_{QA}$  is

$$\begin{aligned} \int_{q_B^+}^1 (q - q_B^+) f(q) dq \int_{q_B^-}^{q_B^+} f(q) dq - \delta_1 &> 0, \\ \int_{q_B^+ + \beta}^1 \beta f(q) dq \int_{q_B^-}^{q_B^+} f(q) dq - \delta_1 &> 0, \\ \beta \bar{\rho} \int_{q_B^-}^{q_B^+} f(q) dq - \delta_1 &> 0, \\ \beta \bar{\rho} \varepsilon - \delta_1 &> 0. \end{aligned}$$

Hence, for any  $\delta_1 \in (0, \min\{(2F^{-1}(\varepsilon) - 1)\rho\varepsilon, \beta\bar{\rho}\varepsilon\})$  we have that  $\sigma_{QQ}\sigma_{AQ} - \delta_1 > \sigma_{AA}\sigma_{QA}$ .

If  $\int_{q_B^-}^{q_B^+} f(q) dq < \varepsilon$  for all  $\varepsilon > 0$  then  $\sigma_{\emptyset_s} \rightarrow 0$ . Since  $\sigma_{QQ}\sigma_{AQ} - \delta_1 > \sigma_{AA}\sigma_{QA}$  implies

$$\lim_{N \rightarrow \infty} \frac{(\sigma_{AA}\sigma_{QA})^{\frac{N-i}{2}}}{(\sigma_{QQ}\sigma_{AQ})^{\frac{N-i}{2}}} \rightarrow 0$$

for  $i = 0, 1$ , we have  $\frac{\pi_t(v, R)}{\pi_t(v, B)} \rightarrow 0$  and, if  $\int_{q_B^-}^{q_B^+} f(q) dq \neq 0$ , also that  $\frac{\pi_R(v, R)}{\pi_R(v, B)} \rightarrow 0$  and  $\frac{\pi_B(v, R)}{\pi_B(v, B)} \rightarrow 0$ .

If  $\int_{q_B^-}^{q_B^+} f(q) dq = 0$  then  $\pi_R(v, R) = \pi_R(v, B) = \pi_B(v, R) = \pi_B(v, B) = 0$ .

On the other hand, if  $\int_{q_B^-}^{q_B^+} f(q) dq \geq \varepsilon$  then there exists a  $\delta_2 > 0$  such that  $\sigma_{\emptyset_s} > \delta_2$ . Define  $\delta = \min\{\delta_1, \delta_2\}$ . By Lemma 2 we have that as  $N$  grows large  $\frac{\pi_t(v, R)}{\pi_t(v, B)} \rightarrow 0$ ,  $\frac{\pi_R(v, R)}{\pi_R(v, B)} \rightarrow 0$  and  $\frac{\pi_B(v, R)}{\pi_B(v, B)} \rightarrow 0$ .

Therefore, equations (2) and (3) then imply  $q_B^- \rightarrow \frac{1}{2}$  and  $q_B^+ \rightarrow \frac{1}{2}$  which in turn implies  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq \rightarrow 0$  and  $\int_{q_B^-}^{q_B^+} f(q) dq \rightarrow 0$ , which contradicts the fact that either  $\int_{\frac{1}{2}}^{q_B^-} f(q) dq \geq \varepsilon$  or  $\int_{q_B^-}^{q_B^+} f(q) dq \geq \varepsilon$  for a fixed  $\varepsilon$ .

Assume now that for all  $\rho > 0$  there exists an  $\bar{N}$  such that for all  $N \geq \bar{N}$ , we have  $\int_{q_B^+}^1 f(q) dq \leq \rho$ . Fix a  $\rho \in (0, \frac{1}{2})$  and the corresponding  $\bar{N}$ . This means that at most a fraction  $\rho$  of voters vote for  $B$  for any  $N \geq \bar{N}$ . In equilibrium of Type 1,  $q_R^+ = \frac{1}{2}$  and all voters who receive signal  $R$  vote for  $R$ . Hence, if a voter is pivotal it must be that at most a fraction  $\rho$  of voters plus one received signal  $R$ . Since  $\rho$  can be chosen as small as desired and  $N$  as large as desired, we have that if a voter is pivotal then the fraction of voters who received signal  $R$  is negligible compared to the fraction of voters who received signal  $B$  and, hence, the probability that the state of nature is  $B$  converges to one when a voter is pivotal

by law of large numbers. By equation (3) this implies  $q_B^+ \rightarrow \frac{1}{2}$  which contradicts the fact that  $\int_{q_B^+}^1 f(q) dq \leq \rho$ .

## Equilibrium of Type 2

We prove next that in an equilibrium of Type 2 we must have  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ . Assume for now that there exists a  $\rho > 0$  such that  $\int_{q_B^+}^1 f(q) dq > \rho$  for all  $N$ . Consider an equilibrium of Type 2 and suppose there exists a  $\varepsilon > 0$  such that  $\int_{q_R^+}^{q_B^+} f(q) dq \geq \varepsilon$  for all  $N$ . We have that a necessary condition for there to be a  $\delta_1 > 0$  such that  $\sigma_{QQ}\sigma_{AQ} - \delta_1 > \sigma_{AA}\sigma_{QA}$  is

$$\begin{aligned} \int_{q_R^+}^1 qf(q) dq \int_{q_B^+}^1 (1-q)f(q) dq - \delta_1 &> \int_{q_R^+}^1 (1-q)f(q) dq \int_{q_B^+}^1 qf(q) dq, \\ \int_{q_R^+}^1 qf(q) dq \int_{q_B^+}^1 f(q) dq - \delta_1 &> \int_{q_R^+}^1 f(q) dq \int_{q_B^+}^1 qf(q) dq, \\ \int_{q_B^+}^1 qf(q) dq \int_{q_R^+}^{q_B^+} f(q) dq - \delta_1 &> \int_{q_B^+}^1 f(q) dq \int_{q_R^+}^{q_B^+} qf(q) dq, \\ \int_{q_B^+}^1 (q - q_B^+)f(q) dq \int_{q_R^+}^{q_B^+} f(q) dq &> \delta_1. \end{aligned}$$

Since  $\int_{q_B^+}^1 f(q) dq > \rho$  for some  $\rho > 0$  then for all  $\bar{\rho} \in (0, \rho)$  there exists a  $\beta > 0$  such that  $\int_{q_B^+ + \beta}^1 f(q) dq > \bar{\rho}$ , fix such  $\rho$  and consider its corresponding  $\beta$ . Moreover, by assumption  $\int_{q_R^+}^{q_B^+} f(q) dq \geq \varepsilon$ . Thus, if we choose any  $\delta_1 \in (0, \beta\hat{\rho}\varepsilon)$  then  $\sigma_{QQ}\sigma_{AQ} - \delta_1 > \sigma_{AA}\sigma_{QA}$ .

Given that  $\int_{q_R^+}^{q_B^+} f(q) dq \geq \varepsilon$  it is true that  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq \geq \varepsilon$  and, hence, there exists a  $\delta_2 > 0$  such that  $\sigma_{\emptyset s} > \delta_2$ . Define  $\delta = \min\{\delta_1, \delta_2\}$ . Then by Lemma 2 we have then that as  $N$  grows large  $\frac{\pi_i(v,R)}{\pi_i(v,B)} \rightarrow 0$ ,  $\frac{\pi_R(v,R)}{\pi_R(v,B)} \rightarrow 0$  and  $\frac{\pi_B(v,R)}{\pi_B(v,B)} \rightarrow 0$ . Equations (2) and (3) then imply  $q_B^+ \rightarrow \frac{1}{2}$  which in turn implies  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ , this contradicts the fact that  $\int_{q_R^+}^{q_B^+} f(q) dq > \varepsilon$  for a fixed  $\varepsilon$ .

Assume now that for all  $\rho > 0$  there exists an  $\bar{N}$  such that for all  $N \geq \bar{N}$ , we have  $\int_{q_B^+}^1 f(q) dq \leq \rho$ . Fix a  $\rho \in (0, \frac{1}{2})$  and the corresponding  $\bar{N}$ . This means that at most a fraction  $\rho$  of voters vote for  $B$  for any  $N \geq \bar{N}$ . In equilibrium of Type 2,  $q_B^- = \frac{1}{2}$  and we have two possibilities. If for all  $\rho > 0$  there exists an  $N$  such that for all  $N \geq \bar{N}$  we have  $\int_{q_R^+}^1 f(q) dq \leq \rho$ , then  $\int_{q_R^+}^{q_B^+} f(q) dq \leq \rho$  which is what the result in the Theorem states. If, on the other hand, there exists a  $\varepsilon > 0$  such that  $\int_{q_R^+}^1 f(q) dq \geq \varepsilon$  for all  $N$ , then at least a fraction  $\varepsilon$  of voters who receive signal  $R$  vote for  $R$ . If a voter is pivotal, it must be because at most a fraction  $\rho$  of voters plus one receive signal  $R$ . However, since  $\rho$  can be chosen as small as desired and  $N + 1$  as large as desired, the fraction of voters who receive signal  $R$  must be arbitrarily small as otherwise a fraction  $\varepsilon$  of them vote for  $R$  against the fraction  $\rho$  that vote for  $B$  and the voter is not pivotal. Therefore, the probability that the state of

nature is  $B$  converges to one when a voter is pivotal by law of large numbers. By equation (3) this implies  $q_B^+ \rightarrow \frac{1}{2}$  which contradicts the fact that  $\int_{q_B^+}^1 f(q) dq \leq \rho$ .  $\square$

*Proof of Proposition 1.* Using the proof of Theorem 2 we have that either  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq \rightarrow 0$  or  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ . Assume first that  $\int_{\frac{1}{2}}^{q_B^+} f(q) dq \rightarrow 0$ . In this case almost all voters vote for the candidate that coincides with their signal (for all  $\delta > 0$  there exists a  $N$  for which the proportion of voters who do not is smaller than  $\delta$ ). Therefore, by law of large numbers the proportion of voters who vote for the candidate that coincides with the state of nature is  $\mu$  while the proportion of voters who vote for the other candidate is  $1 - \mu$ . Since  $\mu > \frac{1}{2}$  implies that there exists a  $\varepsilon > 0$  such that  $\mu - \varepsilon > \frac{1}{2}$ , we have that most voters vote for the candidate that coincides with the state of nature which gives the desired result.

Assume now  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$ . In this case all voters who do not abstain vote for the candidate that coincides with their signal and, furthermore, the decision on whether to vote or not is independent on the signal received (for all  $\delta > 0$  there exists a  $N$  for which the number of voters choose whether to abstain or not depending on their signal is smaller than  $\delta$ ). Therefore, by law of large numbers the proportion of voters who vote for the candidate that coincides with the state of nature is  $(N + 1) \int_{q_B^+}^1 q f(q) dq$  while the proportion of voters who vote for the other candidate is  $(N + 1) \int_{q_B^+}^1 (1 - q) f(q) dq$ . If there exists a  $\rho > 0$  such that  $\int_{q_B^+}^1 f(q) dq > \rho$  for all  $N + 1$  then for all  $\bar{\rho} \in (0, \rho)$  there exists a  $\beta > 0$  such that  $\int_{q_B^+ + \beta}^1 f(q) dq > \bar{\rho}$  we have

$$\begin{aligned} \int_{q_B^+}^1 q f(q) dq - \int_{q_B^+}^1 (1 - q) f(q) dq &= \int_{q_B^+}^1 (2q - 1) f(q) dq \\ &\geq \int_{q_B^+ + \beta}^1 (2q - 1) f(q) dq \\ &\geq 2\beta \int_{q_B^+ + \beta}^1 f(q) dq \\ &> 2\beta \bar{\rho}. \end{aligned}$$

Thus, most voters vote for the candidate that coincides with the state of nature as we wanted to show.

Consider now the case where for all  $\rho > 0$  there exists a  $\bar{N}$  such that  $\int_{q_B^+}^1 f(q) dq \leq \rho$  for all  $n > \bar{N}$ . By monotonicity of  $F$  and the fact that  $\int_{q_R^+}^{q_B^+} f(q) dq \rightarrow 0$  we have  $q_R^+ \rightarrow q_B^+ \rightarrow 1$ .



Moreover,

$$\begin{aligned}
\lim_{q_B^+ \rightarrow 1} \frac{\sigma_{AQ}}{\sigma_{QQ}} &= \lim_{q_B^+ \rightarrow 1} \frac{\int_{q_B^+}^1 (1-q)f(q)dq}{\int_{q_B^+}^1 qf(q)dq}, \\
&= \lim_{q_B^+ \rightarrow 1} \frac{(1-q_B^+)f(q_B^+)}{q_B^+f(q_B^+)}, \\
&= \lim_{q_B^+ \rightarrow 1} \frac{1}{q_B^+} - 1, \\
&= 0,
\end{aligned}$$

and similarly  $\lim_{q_R^+ \rightarrow 1} \frac{\sigma_{QA}}{\sigma_{AA}} = 0$ , where we have used L'Hôpital's rule for computing the limit above. That is, the probability that a random voter votes for the candidate that does not match the state of nature is insignificant compared to the probability that a random voter votes for the candidate that does, which implies  $P(V = S) \rightarrow 1$ .  $\square$

## B Appendix: Experimental Instructions

Welcome! You are about to participate in a decision making experiment. If you follow the instructions carefully, you can earn a considerable amount of money depending on your decisions and the decisions of the other participants. Your earnings will be paid to you in cash at the end of the experiment

This set of instructions is for your private use only. During the experiment you are not allowed to communicate with anybody. In case of questions, please raise your hand. Then we will come to your seat and answer your questions. Any violation of this rule excludes you immediately from the experiment and all payments.

For your participation you will receive a show-up fee 3 pounds. You can earn additional amounts of money. Below we will describe how. All your decisions will be treated confidentially both during the experiment and after the experiment. This means that none of the other participants will know which decisions you made.

**Experimental Instructions** The experiment will last for 30 rounds. In each round you will be matched randomly in groups of four participants. Remember that the groups change in each round, so the participants you play with in one round are most likely different from those you played with the round before. At the beginning of each round of the experiment the computer randomly draws one of two colours RED or BLUE. We call the colour that was drawn “the state”. BLUE is much more likely than RED to be drawn. In particular there is a 95% chance that BLUE is drawn and only a 5% chance that RED is drawn. *Remember that the state is drawn anew in each round, i.e. it can be different in each round. The state is the same, though, for all group members in each round.*




Figure 7: The state is BLUE with a 95% chance, i.e. a chance of 95 in 100.

**Goal of the experiment:** You will be asked to guess whether the state is RED or BLUE. Your goal is to guess correctly as a group. Hence it will not matter whether you guess correctly yourself. The only thing that matters is whether the majority of your group guesses correctly. We will explain now what additional information each group member gets before making a guess, what guesses you can make and how your payments are computed.

**Information you receive:** Each group member receives a “signal” about whether the state is BLUE or RED before they submit their guess. A signal is a ball drawn randomly from a box containing RED and BLUE balls. All balls in a box are equally likely to be drawn.

There are however, two boxes for each player and you don’t know which one the ball is drawn from. If the state is BLUE the ball will be drawn from your BLUE box. (Remember that this is the case with a 95% chance). If the state is RED, the ball will be drawn from your RED box. (This is the case with a 5% chance). Hence if you knew the box you would know the state. This is true for all participants.

There are always at least as many BLUE balls in your BLUE box as there are in your RED box. Hence, if both boxes were equally likely, a BLUE ball is more likely to come from a BLUE box and a RED ball is more likely to come from a RED box.

How much more likely will depend on the exact composition of the boxes. In each round you will be shown the composition of your boxes. You will also be shown the colour of the ball drawn.

It is important to note that the composition of boxes can be different for different group members. In particular, for each participant, the number of BLUE balls in their BLUE box is randomly drawn from anything between half the balls being BLUE to all balls being BLUE. The number of RED balls in a participant’s RED box always equals the number of BLUE balls in their BLUE box

Things to remember about signals:

- You will see a ball drawn from either your RED or your BLUE box.
- If the state is BLUE the ball will be drawn from the BLUE box. If the state is RED it will be drawn from the RED box.
- You will also see how many RED and BLUE balls your RED and BLUE boxes contain.
- All other group members will also see a ball drawn from one of their boxes.
- Remember, though, that their boxes can have a different composition.
- Boxes change in each round for each participant.

**Making a guess:** After all group members have received their signals, all will make a guess simultaneously. You have three options. You either guess RED, BLUE or you can ABSTAIN. Remember that the goal is to guess correctly as a group.

**Your payment:** Apart from the show up fee you receive, one round is drawn for payment and you receive

- 10 additional pounds if the group guesses correctly in that round and
- 2 additional pounds if the group is not correct in that round.

**When is the group correct?** The group is correct if the majority of group members who do not abstain indicate the correct state.

Hence, if the state is BLUE then the group is correct if

- at least 3 group members vote BLUE,
- at least 2 group members vote BLUE and at least one abstains,
- at least 1 group member votes BLUE and all others abstain.

Similarly, if the state is RED then the group is correct if

- at least 3 group members vote RED,
- at least 2 group members vote RED and at least one abstains,
- at least 1 group member votes RED and all others abstain.

If the same number of group members vote RED and BLUE, then there is a tie and whether the group's guess is considered correct is determined by the flip of a coin.

**Control Questions:** Are the following statements TRUE or FALSE? If you have any questions please raise your hand.

1. My group members change from round to round.
2. All group members receive a ball from the same box.
3. The composition of the box of my group members can be different from the composition of my box.

4. If I vote RED, one group member abstains and two vote BLUE, I receive 2 pounds if the state is RED and 10 pounds if the state is BLUE.
5. If I vote BLUE, one group member abstains and two vote BLUE, I receive 2 pounds if the state is RED and 10 pounds if the state is BLUE.
6. Only one round is randomly drawn for payment.

**ENJOY THE EXPERIMENT !!**

## C Appendix: Additional Tables

This Appendix collects additional tables. Table 6 shows the distribution of votes with/against the signal and abstentions depending on treatment and signal. In the asymmetric treatments participants usually vote with their signal if it is BLUE. If their signal is RED, by contrast, the modal action is to vote against the signal. There is a substantial share of abstentions as well among voters with a RED signal.

	SYM			ASYM			ASYM-COARSE		
	RED	BLUE	Overall	RED	BLUE	Overall	RED	BLUE	Overall
with signal	0.65	0.72	0.69	0.19	0.90	0.66	0.09	0.89	0.69
abstain	0.23	0.21	0.22	0.34	0.08	0.17	0.24	0.10	0.17
against signal	0.12	0.07	0.09	0.46	0.02	0.17	0.67	0.01	0.17

Table 6: Share of votes with/against the signal as well as abstentions depending on treatment and signal.

	(1)	(2)	(3)	(4)
Vote against signal if it is:	RED	BLUE	RED 15-30	BLUE 15-30
q	-0.0698 (0.122)	-0.243*** (0.0586)	-0.149 (0.147)	-0.192** (0.0864)
<b>ASYM</b>	0.434** (0.171)	-0.172*** (0.0600)	0.547*** (0.200)	-0.183** (0.0846)
<b>ASYM-COARSE</b>	0.900*** (0.239)	-0.192** (0.0751)	0.688** (0.290)	-0.184* (0.104)
<b>ASYM</b> × q	-0.137 (0.223)	0.178** (0.0789)	-0.267 (0.261)	0.153 (0.114)
<b>ASYM-COARSE</b> × q	-0.619* (0.369)	0.171 (0.112)	-0.191 (0.461)	0.138 (0.155)
Constant	0.169 (0.105)	0.243*** (0.0434)	0.194 (0.125)	0.224*** (0.0619)
Observations	809	1,351	435	645
Groups	72	72	72	72

Robust standard errors in parentheses

\*\*\* p<0.01, \*\* p<0.05, \* p<0.1

Table 7: Random Effects OLS regressions: Voting against one's signal when it indicates the low prior (column (1)) or high prior (column (2)) state. Columns (3) and (4) only consider data from the last 15 periods.

$$y_{it} = \alpha_i + \beta_0 q_{it} + \beta_1 \mathbf{ASYM} + \beta_2 \mathbf{ASYM-COARSE} + \beta_{10} * (q_{it} * \mathbf{ASYM}) + \beta_{20} * (q_{it} * \mathbf{ASYM-COARSE}) + \epsilon_{it} \quad (11)$$

Table 7 shows the results of running regression (11) in our sample using as binary outcome  $y_{it}$  whether or not a participant  $i$  abstained in period  $t$ . Columns (1) and (2) include the whole sample, columns (3) and (4) only the second half of the experiment after potentially some learning has occurred. Columns (1) and (3) focus on participants who received a RED signal. The estimates show that in the baseline (treatment **SYM**) participants rarely vote against their signal. They do so more often in **ASYM** and **ASYM-COARSE** if the signal is RED, i.e. goes against the prior and less often if it is BLUE, i.e. consistent with the prior. Signal accuracy decreases the propensity to vote against the signal across all treatments, albeit not always significantly so.

	(1)	(2)	(3)	(4)
Abstain if signal is	RED	BLUE	RED 15-30	BLUE 15-30
q	-1.171*** (0.125)	-1.148*** (0.106)	-1.258*** (0.169)	-1.174*** (0.143)
<b>ASYM</b>	-0.396** (0.173)	-0.527*** (0.112)	-0.610*** (0.223)	-0.697*** (0.145)
<b>ASYM-COARSE</b>	-1.065*** (0.242)	-0.731*** (0.138)	-1.004*** (0.328)	-0.890*** (0.175)
<b>ASYM</b> × q	0.636*** (0.228)	0.627*** (0.142)	1.064*** (0.301)	0.751*** (0.188)
<b>ASYM-COARSE</b> × q	1.600*** (0.378)	0.896*** (0.201)	1.528*** (0.528)	1.061*** (0.257)
Constant	1.060*** (0.105)	0.997*** (0.0811)	1.123*** (0.138)	1.056*** (0.106)
Observations	809	1,351	435	645
Number of id	72	72	72	72

Robust standard errors in parentheses

\*\*\* p<0.01, \*\* p<0.05, \* p<0.1

Table 8: Random Effects OLS regressions: Abstaining when the signal indicates the low prior (column (1)) or high prior (column (2)) state. Columns (3) and (4) only consider data from the last 15 periods.

Table 8 shows the results of running regression (11) in our sample using as binary outcome  $y_{it}$  whether or not a participant  $i$  abstained in period  $t$ . Irrespective of the signal, participants abstain less often in **ASYM** and **ASYM-COARSE** compared to **SYM**. An increased

signal accuracy increases the propensity to abstain in **ASYM** and **ASYM-COARSE**, where participants decide to abstain rather than voting against the signal, but decreases it in **SYM**, where participants decide to vote with the signal instead of abstaining.