

Group Theory: Math30038, Sheet 2

GCS

The course web site is available via <http://www.bath.ac.uk/~masgcs/>

1. Let G be a finite group of even order. Show that G must contain an element of order 2.
2. Suppose that G is a cyclic group and that $|G| = 12$. Make a list of all the subgroups of G , and draw a “Hasse diagram”. This is a diagram of dots and lines (as shown in a lecture), where the inclusion $A \leq B$ is indicated by a line or sequence of lines going up from the dot labelled A to the dot labelled B . You can label the lines with corresponding indices. Stare at your diagram, and think about the following set: $\{d \in \mathbb{N} \mid d \text{ divides } 12\}$.
3. Suppose that G is a group and that $|G| = 4$. By considering the orders of the elements of G (or otherwise), prove that G must be abelian.
4. Suppose that G is a group and that $|G| = 6$.
 - (a) Show that if G is abelian, then G must be cyclic.
 - (b) Show that if G is nonabelian, then G must contain an element x of order 2 and an element y of order 3, and moreover it must be the case that $xyx = y^{-1}$.
5. Consider a cube in Euclidean space. The group R of all rigid motions of the cube (reflections not allowed) has order 24. Paint the vertices black and white so that if two vertices are joined by an edge, then they have opposite colour. Let G be the subgroup of R consisting of those rigid motions which preserve the colouring (black corners go to black corners and white corners to white ones). Describe the 12 elements of G . Show that G has no subgroup of order 6.
6. Let $\Omega = \mathbb{Z}$. Let $\text{Sym}(\Omega)$ denote the group of all bijections from Ω to Ω under composition of maps. Let $\alpha \in \text{Sym}(\Omega)$ be such that $(x)\alpha = x + 2 \forall x \in \mathbb{Z}$ and let β be the bijection which swaps 0 and 1, but fixes all other integers. Let $G = \langle \alpha, \beta \rangle$, so G is a finitely generated subgroup of $\text{Sym}(\Omega)$. Let $X = \{\alpha^{-i}\beta\alpha^i \mid i \in \mathbb{Z}\}$. Let $H = \langle X \rangle$. Show that H is not finitely generated (i.e. there does not exist a finite subset Y of H such that $H = \langle Y \rangle$). *Hint: the support of a bijection is the set of elements which are not fixed by the bijection. Show that all elements of H have finite support, and think about the consequences of this.*

7. Consider the group of bijections from \mathbb{R}^3 to \mathbb{R}^3 under composition of maps. Let i denote the bijection which is clockwise rotation through $\pi/2$ about the x axis, and j denote the bijection which is clockwise rotation through $\pi/2$ about the y axis. Let $k = ij$. Describe k geometrically. Discuss the group $\langle i, j \rangle$: what is its order and what does its multiplication table look like?